



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification⁴ : C12N 9/22, 5/02, 15/00 C12N 1/20, C12P 19/34 C07H 21/02, A61K 31/73, 39/235 A61K 39/285</p>	<p>A1</p>	<p>(11) International Publication Number: WO 89/ 05852 (43) International Publication Date: 29 June 1989 (29.06.89)</p>
<p>(21) International Application Number: PCT/AU88/00478 (22) International Filing Date: 14 December 1988 (14.12.88) (31) Priority Application Numbers: PI 5911 PI 9950 PJ 0353 PJ 1304 PJ 1333 (32) Priority Dates: 15 December 1987 (15.12.87) 19 August 1988 (19.08.88) 9 September 1988 (09.09.88) 4 November 1988 (04.11.88) 7 November 1988 (07.11.88) (33) Priority Country: AU (71) Applicant (for all designated States except US): COMMON-WEALTH SCIENTIFIC AND INDUSTRIAL RE-SEARCH ORGANISATION [AU/AU]; Limestone Avenue, Campbell, ACT 2601 (AU). (72) Inventors; and (75) Inventors/Applicants (for US only) : HASELOFF, James, Phillip [AU/AU]; 17 Hemmant Street, O'Connor, ACT 2601 (AU). GERLACH, Wayne, Lyle [AU/AU]; 22 Goble Street, Hughes, ACT 2605 (AU). JENNINGS, Philip, Anthony [GB/AU];</p>		<p>13 Lobelia Street, West Chatswood, NSW 2066 (AU). CAM-ERON, Fiona, Helen [AU/AU]; 2 Rounce Avenue, Forest-ville, NSW 2087 (AU). (74) Agents: CORBETT, Terence, Guy et al.; Davies & Collison, 1 Little Collins Street, Melbourne, VIC 3000 (AU). (81) Designated States: AT, AU, BB, BG, BJ (OAPI patent), BR, CF (OAPI patent), CG (OAPI patent), CH, CM (OAPI pa-tent), DE, DK, FI, GA (OAPI patent), GB, HU, JP, KP, KR, LK, LU, MC, MG, ML (OAPI patent), MR (OAPI pa-tent), MW, NL, NO, RO, SD, SE, SN (OAPI patent), SU, TD (OAPI patent), TG (OAPI patent), US. Published <i>With a supplementary international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> Date of publication of the supplementary international search report: 25 January 1990 (25.01.90)</p>
<p>(54) Title: RIBOZYMES</p> <p style="text-align: center;">AA G C*G AGXAGUC N₁*N₂ (I)</p>		
<p>(57) Abstract</p> <p>Synthetic single stranded RNA molecule(s) and derivatives thereof possessing highly specific endoribonuclease activity, hereinafter defined as ribozyme. The ribozyme cleaves target RNA sequence by hybridizing partially to the target RNA. It comprises of a hybridizing region and a catalytic region. The hybridizing region contains one or more arms of single stranded RNA. The preferred ribozyme has a catalytic region consisting of the sequence (I), in a loop structure where * represents base pairing between complementary nucleotides, X is any ribonucleotide and N₁ and N₂ represent ribonucleo-tides of complementary sequences along at least part of the length. The ribozyme cleaves target RNA after the triplet XUY, where X is any ribonucleotide, U is uracil and Y is adenine, cytosine or uracil. Methods for inactivating RNA (in vitro or in vivo), methods of preparing the ribozymes, and compositions containing the ribozymes are also described.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MR	Mauritania
BE	Belgium	GA	Gabon	MW	Malawi
BF	Burkina Faso	GB	United Kingdom	NL	Netherlands
BG	Bulgaria	HU	Hungary	NO	Norway
BJ	Benin	IT	Italy	RO	Romania
BR	Brazil	JP	Japan	SD	Sudan
CA	Canada	KP	Democratic People's Republic of Korea	SE	Sweden
CF	Central African Republic	KR	Republic of Korea	SN	Senegal
CG	Congo	LI	Liechtenstein	SU	Soviet Union
CH	Switzerland	LK	Sri Lanka	TD	Chad
CM	Cameroon	LU	Luxembourg	TG	Togo
DE	Germany, Federal Republic of	MC	Monaco	US	United States of America
DK	Denmark				

RIBOZYMES

The present invention relates to a class of synthetic RNA molecules and derivatives thereof, hereinafter referred to as ribozymes, which possess highly specific endoribonuclease activity.

5 A number of naturally occurring RNA molecules such as avocado sunblotch viroid (ASBV), the satellite RNAs of tobacco ringspot virus (sTobRV) and lucerne transient streak virus (sLTSV), undergo self-catalysed cleavage. Such cleavage appears to be
10 an essential and unique part of the life cycle of these and other RNAs.

Self-catalysed RNA cleavage reactions share a common requirement for divalent metal ions and neutral or higher pH, and result in the production of
15 RNA with termini possessing 5' hydroxyl and 2',3' cyclic phosphate groups (Prody et al., Science 231: 1577-1580 (1986) and Buzayan et al., Virology 151: 186-199 (1986)). The cleavage reactions are catalysed by the RNAs themselves, presumably as a

result of conformation bringing reactive groups into close proximity. The sites of self-catalysed cleavage in naturally occurring RNAs are located within highly conserved regions of RNA secondary structure (Buzayan et al., Proc. Natl. Acad. Sci. USA 83: 8859-8862 (1986) and Forster, A.C. and Symons, R.H. Cell 50: 9-16 (1987)).

Experiments we have carried out on the satellite RNAs of tobacco ringspot virus (sTobRV) have led to the design of novel endoribonucleases (hereinafter referred to as "ribozymes"), that is, enzymes comprised of RNA which catalyse specific cleavage of RNA target molecules.

The term ribozyme as used in the specification refers to molecules comprised wholly of RNA or derivatives thereof.

The ribozymes of the present invention are distinct from the RNA endoribonuclease which occurs naturally in Tetrahymena Thermophila (known as the IVS, or L-19 IVS RNA) and which has been extensively described by Thomas Cech and collaborators (Zaug, A.J. et al, Science (1984) 224: 574-578; Zaug, A.J. and Cech, T.R., Science (1986) 231: 470-475; Zaug, A.J., et al, Nature (1986) 324: 429-433; published International patent application No. WO 88/04300 by University Patents Inc.). The Cech endoribonuclease has an eight base pair active site which hybridizes to a target RNA sequence whereafter cleavage of the target RNA takes place, with a requirement for free guanosine or guanosine derivatives. The fragments which arise from cleavage contain terminal 5' phosphate and 3' hydroxyl groups. The limited number of nucleotides available for hybridization to an RNA substrate limits the effectiveness or efficiency of

the Cech endoribonuclease as in general, oligonucleotides comprising less than twelve nucleotides hybridize poorly to target sequences. It also appears that a number of nucleotides in the active site of the Cech endoribonuclease may need to be conserved for efficient endoribonuclease activity. This restricts the number of permutations of active site sequences which can be engineered to effect hybridization to target sequences, thereby restricting the range of RNA target sequences cleavable by the Cech endoribonuclease. The Cech endoribonuclease also modifies RNA by adding a free guanosine nucleotide to the 5' end of cleaved RNA.


In contrast, the ribozymes of the present invention hybridize efficiently to a wide range of target RNA sequences, and do not modify the cleaved target RNA.

The ribozymes of the present invention comprise a hybridizing region which is complementary in nucleotide sequence to at least part of a target RNA, and a catalytic region which is adapted to cleave the target RNA. The hybridizing region contains 9 or more nucleotides.

In a preferred aspect the ribozymes of the present invention have a hybridizing region comprising one or more arms formed of single stranded RNA and having a sequence complementary to at least part of a target RNA, said one or more arms being associated with a catalytic region capable of cleaving said target RNA; and where the hybridizing region comprises a single arm of RNA said arm contains at least 9 nucleotides, and where the hybridizing region comprises 2 or more arms of RNA, the sum of nucleotides in said arms is greater than 9

nucleotides.

In one embodiment of the invention, there is provided a ribozyme of the formula 1:



wherein; X represents any ribonucleotide and each X
 15 residue may be the same or different;
 the sum of n and n' is greater than 6 and n
 and n' may be the same or different;
 an (*) represents a base pair between
 complementary ribonucleotides;
 20 X' and X'' represent oligoribonucleotides of
 complementary sequence along at least part
 of their length to allow base pairing
 between the oligoribonucleotides, or X' and
 X'' together form a single RNA sequence
 25 wherein at least part of said sequence
 comprises a stem formed by base pairing
 between complementary nucleotides; and
 optionally, an additional nucleotide
 selected from one of A, G, C or U may be
 30 inserted after ¹A in formula (1).


Region (I) of formula (1) represents the arms or
 flanking sequences of a ribozyme which hybridize to
 respective portions of a target RNA sequence. The
 arms may hybridize along the full length of the

target RNA or part thereof. An RNA catalytic region is depicted at region (II) of formula 1. The catalytic region may contain one or more additional nucleotides which do not adversely effect catalytic activity. Such additions could be readily tested for ribozyme activity without undue experimentation following the teachings of the specification. The catalytic region may also form part of the hybridizing region.

10 The oligoribonucleotides X' and X" may comprise up to 5,000 or more nucleotides.

According to a further specific embodiment of the present invention there is provided a ribozyme of the formula 2:

15



wherein; X represents any ribonucleotide and each X residue may be the same or different;

30 an (*) represents a base pair between complementary ribonucleotides;

n and n' are as previously defined;

m and m' are 1 or more and may be the same or different;

B represents a bond, a base pair, a ribonucleotide, or an oligoribonucleotide containing at least 2 ribonucleotides;

and optionally, an additional nucleotide
5 selected from any one of A, G, C or U may be inserted after ¹A in formula (2).

The ribozymes of the present invention may be prepared by methods known per se in the art for the synthesis of RNA molecules. (For example, according
10 to recommended protocols of Promega, Madison, WI, USA). In particular, the ribozymes of the invention may be prepared from a corresponding DNA sequence (DNA which on transcription yields a ribozyme, and which may be synthesized according to methods known
15 per se in the art for the synthesis of DNA) operably linked to an RNA polymerase promoter such as a promoter for T7 RNA polymerase or SP6 RNA polymerase. A DNA sequence corresponding to a ribozyme of the present invention may be ligated into
20 a DNA transfer vector, such as plasmid or bacteriophage DNA. Where the transfer vector contains an RNA polymerase promoter operably linked to DNA corresponding to a ribozyme, the ribozyme may be conveniently produced upon incubation with an RNA
25 polymerase. Ribozymes may, therefore, be produced in vitro by incubation of RNA polymerase with an RNA polymerase promoter operably linked to DNA corresponding to a ribozyme, in the presence of ribonucleotides. In vivo, prokaryotic or eukaryotic
30 cells (including mammalian and plant cells) may be transfected with an appropriate transfer vector containing genetic material corresponding to a ribozyme in accordance with the present invention, operably linked to an RNA polymerase promoter such

that the ribozyme is transcribed in the host cell. Transfer vectors may be bacterial plasmids or viral RNA or DNA. Nucleotide sequences corresponding to ribozymes are generally placed under the control of strong promoters such as, for example, the lac, SV40 late, SV40 early, metallothionin, or λ promoters. Ribozymes may be directly transcribed in-vivo from a transfer vector, or alternatively, may be transcribed as part of a larger RNA molecule. For example, DNA corresponding to ribozyme sequences may be ligated into the 3' end of a carrier gene, for example, after a translation stop signal. Larger RNA molecules may help to stabilize the ribozyme molecules against nuclease digestion within cells. On translation the carrier gene may give rise to a protein, whose presence can be directly assayed, for example, by enzymatic reaction. The carrier gene may, for example, encode an enzyme.

In a further aspect of the invention, there is provided a DNA transfer vector which contains a DNA sequence corresponding to a ribozyme operably linked to a promoter to provide transcription of the ribozyme.

In one preferred method of producing a ribozyme, two synthetic oligonucleotides of complementary sequence are prepared by standard procedures (for example, using an Applied Biosystems Model 380A DNA Synthesizer (Applied Biosystems Inc., Foster City, California 94404)), and hybridized together. One of the oligonucleotides encodes a desired ribozyme. The respective ends of the hybridized oligonucleotides correspond to different restriction enzyme sites, say EcoRI at one end and PstI at the other end. After cleavage with appropriate restriction enzymes (EcoRI

and PstI in the above example), the double stranded DNA fragment may be cloned into a transfer vector. Where the plasmid vector contains an RNA polymerase promoter upstream from the DNA sequence corresponding
5 to a ribozyme of the present invention, RNA transcripts corresponding to the ribozyme can be conveniently prepared either in-vitro or in-vivo. Where the ribozyme is comprised of two halves held together by base-pairing between complementary
10 nucleotides, each half of the ribozyme may be produced according to the above methods, and the halves incubated together to form the ribozyme.

The preferred ribozymes of the present invention cleave target RNA which contains the sequence X^OUY ,
15 where X^O is any ribonucleotide, U is uracil and Y is adenine, cytosine or uracil. X^OU forms part of a base pair flanking region and Y is not base paired. Preferably, but by no means exclusively, X^O is guanidine, and X^OUY is GUC or GUA. Any RNA
20 molecule containing these sequences can be cleaved with the ribozymes of the present invention. Once the sequence of an RNA transcript containing the sequence X^OUY has been determined, the arms of the ribozyme sequence can be synthesised to be
25 complementary to, and thus hybridizable to, the RNA on the target sequence flanking the X^OUY sequence. On hybridization of the arms of the ribozyme to the target RNA sequence flanking the X^OUY sequence, the catalytic region of the ribozyme cleaves the target
30 RNA within the X^OUY sequence. RNA cleavage is facilitated in the presence of magnesium or other divalent cation at a pH of approximately 8.0.

Accordingly, the preferred ribozymes of the present invention can be engineered to cleave any RNA

whose sequence is known. The high frequency of the residues cleaved by the ribozymes in RNA (1:64 for GUC in an RNA with random and equal frequency of base distribution) means that a number of potential sites
5 for ribozyme cleavage can be confidently predicted in any given target RNA.

According to another aspect of the present invention there is provided a method for the inactivation of a target RNA sequence, which
10 comprises reacting said target RNA sequence with a ribozyme of the present invention.

In-vivo, that is, within the cell or cells of an organism, a transfer vector such as a bacterial plasmid or viral RNA or DNA, encoding one or more
15 ribozymes, may be transfected into cells e.g. (Llewellyn et al., J. Mol. Biol. (1987) 195: 115-123; Hanahan et al. J. Mol. Biol (1983) 166). Once inside the cell, the transfer vector may replicate, and be transcribed by cellular polymerases
20 to produce ribozyme RNAs which then inactivate a desired target RNA. Alternatively, a transfer vector containing one or more ribozyme sequences may be transfected into cells or introduced into cells by way of micromanipulation techniques such as
25 microinjection, such that the transfer vector or a part thereof becomes integrated into the genome of the host cell. Transcription of the integrated genetic material gives rise to ribozymes, which act to inactivate a desired target RNA.

30 The ribozymes of the present invention have extensive therapeutic and biological applications. For example, disease causing viruses in man and animals may be inactivated by administering to a subject infected with a virus, a ribozyme in

accordance with the present invention adapted to hybridize to and cleave RNA transcripts of the virus. Such ribozymes may be delivered by parenteral or other means of administration. Alternatively, a
5 subject infected with a disease causing virus may be administered a non-virulent virus such as vaccinia or adenovirus which has been genetically engineered to contain DNA corresponding to a ribozyme operably linked to an RNA promoter, such that the ribozyme is
10 transcribed in the cells of the host animal, transfected with the engineered virus, to effect cleavage and/or inactivation of the target RNA transcript of the disease causing virus. The ribozymes of the present invention have particular
15 application to viral diseases caused for example, by the herpes simplex virus (HSV) or the AIDS virus (HIV).

The ribozymes of the present invention also have particular application to the inactivation of RNA
20 transcripts in bacteria and other prokaryotic cells, plants and animals. In bacteria, RNA transcripts of, for example, bacteriophage, which cause bacterial cell death, may be inactivated by transfecting a cell with a DNA transfer vector which is capable of producing a
25 ribozyme in accordance with the present invention which inactivates the phage DNA. Alternatively, the ribozyme itself may be added to and taken up by the bacterial cell to effect cleavage of the phage RNA.

RNA transcripts in plants may be inactivated
30 using ribozymes encoded by a transfer vector such as the Ti plasmid of Agrobacterium tumefaciens. When such vectors are transfected into a plant cell, the ribozymes are produced under the action of RNA polymerase and may effect cleavage of a specific

target RNA sequence. Accordingly, plant viruses whose RNA sequence are known, or the RNA transcripts of plant genes, may be inactivated using ribozymes.

Endogenous gene transcripts in plants, animals or
5 other cell types may be inactivated using the ribozymes of the present invention. Accordingly, undesirable phenotypes or characteristics may be modulated. It may, for example, be possible using the ribozymes of the present invention to remove
10 stones from fruit or treat hereditary diseases in humans which are caused by the production of a deleterious protein, or over production of a particular protein.

The present invention will now be illustrated by
15 way of non-limiting example only, with reference to the following non-limiting Examples, and Figures.

In the Figures:

FIGURE 1 shows RNA self cleavage sites of wild type and mutated RNAs; and an electrophoretic profile
20 showing self-catalysed RNA cleavage products.

(a) Summarises the conserved structures associated with naturally-occurring RNA cleavage sites in ASBV, newt satellite DNA transcripts and the satellite RNAs of sTobRV, LTSV, velvet tobacco mottle
25 virus, solanum nodiflorum mottle virus and subterranean clover mottle virus. Nucleotide sequences which are conserved between these structures are shown, while others are represented as X. Base-pairing is represented by "*" and the site
30 for RNA cleavage is arrowed.

(b) Shows the conserved nucleotide sequence associated with the cleavage of the (+) strand of sTobRV RNA. The cleavage site is arrowed.

(c) An in vitro mutant of sTobRV containing an

insertion of eight nucleotides (shown boxed) together with a flanking duplication of three nucleotides (UGU residues 7 to 9) is shown.

(d) Sub-cloned HaeIII fragments of wild type
5 sTobRV and the D-51 in vitro mutant were each transcribed in both the (+) and (-) orientations and radiolabelled transcripts fractionated by polyacrylamide gel electrophoresis. The positions of uncleaved 159 and 170 base transcripts from the wild
10 type (WT) and mutant (D-51) sequences are arrowed; sizes of cleavage products are shown.

FIGURE 2 shows the nucleotide sequence of a ribozyme and the products of ribozyme cleavage separated by gel electrophoresis.

15 (a) The inserted nucleotides in the D-51 mutant (FIG. 1c) contains a BamHI restriction endonuclease site. BamHI was used to split the mutant DNA, and the two sequences were sub-cloned and transcribed separately in vitro. The RNA transcripts are shown
20 schematically, with potential base-pairings between the RNAs indicated "**". The fragment containing the arrowed site for cleavage is referred to as S-RNA, the fragment containing the ribozyme is designated Rz-RNA.

25 (b) [32 P]-Rz-RNA (101 bases) was incubated alone (lane 1), and with unlabelled S-RNA (lane 2). [32 P]-S-RNA was incubated alone (lane 3), and with unlabelled and 32 P labelled Rz-RNAs (lanes 4 & 5 respectively).

30 FIGURE 3 shows a schematic model of a ribozyme according to one embodiment of the present invention. Region A represents the cleavage sequence within the target RNA. Region B represents a catalytic region, and regions C represent the arms of

the ribozyme.

FIGURE 4 shows the design of ribozymes targeted against the CAT (chloramphenicol acetyl transferase) gene transcript. Ribozymes, termed RzCAT-1, 2 and 3, were targeted against three sites within an 835 base in vitro transcript of the CAT gene. The relative locations of the cleavage sites on the transcript are shown schematically with the flanking bases numbered (a). The three ribozyme sequences are shown ((b) to (d)) with their target sequences. Amino-acid sequences of the CAT gene are numbered and the predicted sites for RNA cleavage arrowed. RzCAT-1 and 3 contain 24 base sequences derived from (+) strand sTobRV (region B, FIG. 3), while RzCAT-2 contains a single U-A change in this region.

FIGURE 5 shows the results of CAT RNA cleavage with ribozymes RzCAT-1 to 3.

(a) The [32 P]-CAT RNAs were gel fractionated after incubation alone (-) or with with one of the three ribozymes, RzCAT-1 to 3 (lanes 1, 2 & 3 respectively). The location of full-length transcript is shown arrowed.

(b) 5' Terminal base analysis. The 3' fragments produced by ribozyme cleavage of CAT mRNA were [32 P]-kinased, gel purified, subjected to complete nuclease digestion and the released terminal residues fractionated by pH 3.5 polyacrylamide gel electrophoresis. The 5' terminal nucleotides, determined by reference to markers (lane M), were A, U and G for the fragments produced by RzCAT-1 to 3 (lanes 1, 2 and 3 respectively).

FIGURE 6 shows a time course of ribozyme (RzCAT-1) catalytic activity against CAT RNA. The amounts of the 139 nucleotide cleavage product were

quantified and plotted. The inset shows the accumulation of the 139 base fragment with time, after polyacrylamide gel electrophoresis.

FIGURE 7 shows the relative rates of cleavage of CAT RNA under different temperature conditions. Substrate RNA is represented by a solid line. In each case cleavage product is represented by a broken line.

FIGURE 8 depicts three ribozymes (corresponding to RzCAT-2) having arms or flanking sequence of varying length.

FIGURE 9 shows the scheme for production of a ribozyme comprising catalytic anti-sense RNA containing each of the cleavage domains RzCAT-1 to 3.

FIGURE 10 shows ribozymes hybridizing to target sequences containing the GUA (10a) and GUU (10b) motifs in CAT mRNA.

FIGURE 11 shows sites for self catalysed RNA cleavage in citrus exocortis viroid (CEV) RNA and its complement.

FIGURE 12 shows the ribozyme RzCEV25x(+) hybridized to target CEV RNA(a), and a gel electrophoretic profile of (+)CEV RNA and complementary (-) CEV RNA incubated with RzCEV25x(+) (b, lanes 1 and 2 respectively. Cleavage product is arrowed).

FIGURE 13 shows the ribozyme RzCAT-2 hybridizing to its target sequence (a) and ribozyme RzSCMoV(b). The catalytic domain in each ribozyme is boxed. Differences in the catalytic region of RzSCMoV, when compared with RzCAT-2 are marked.

FIGURE 14 shows the ribozyme RzCEV-2 hybridizing to a target sequence in citrus exocortis viroid (CEV) RNA. The cleavage site corresponds to nucleotide-336

in the CEV RNA sequence. The alteration in nucleotide sequence in the catalytic domain, when compared with the catalytic domain of sTobRV, is circled (a). Fig 14(b) shows an electrophoretic
5 profile of a control [(-) strand of CEV] RNA, lane 7 and the (+) strand of CEV RNA lane 8, after incubation with RzCEV2.

FIGURE 15 shows the ribozyme RzCAT-2 (a) compared with ribozyme RzCAT-2B (b). Catalytic domains are
10 boxed. Changes in the catalytic domain of RzCAT-2B compared with RzCAT-2 are also boxed;

FIGURE 16 shows a map of plasmid pJ35SN; and

FIGURE 17 is a graphic presentation of the average of four experiments on the inhibition of CAT
15 expression in plants (tobacco protoplasts).

The following Examples are given to illustrate the present invention and are not to be construed as limiting the present invention.

Reactions and manipulations involving DNA, such
20 as ligations, restriction enzyme digestions, bacterial transformation, DNA sequencing etc. were carried out according to standard techniques, such as those described by Maniatis et al (Molecular Cloning, Cold Spring Harbour, 1982). Manipulations involving
25 RNA were also carried out according to standard techniques, such as those described by Uhlenbeck (Nature 328, 596-600 (1987)) and Haseloff and Gerlach (Nature 334, 585-591 (1988)).

30 EXAMPLE 1

Self-Catalyzed Cleavage of Mutated sTobRV RNA:

A consensus of the domains associated with naturally-occurring RNA cleavage sites in ASBV, newt satellite DNA transcripts and the satellite RNAs of

sTobRV, LTSV, velvet tobacco mottle virus (VMoV), solanum nodiflorum mottle virus (SNMV) and subterranean clover mottle virus (SCMoV) as shown in Figure 1a. Nucleotide sequences which are conserved between these structures are shown, while non-conserved sequences are represented as X. An extra U is positioned after residue ¹A in LTSV(+) strand.

The domain associated with the self-catalysed cleavage of the (+) strand of sTobRV was studied to ascertain the enzymic substrate activity within this domain. First, cloned sTobRV cDNAs were mutagenised using an oligonucleotide linker (BamHI) insertion protocol.

Construction of a Vector for in vitro Expression of sTobRV:

A 160 bp Taq I - Spe I fragment of sTobRV cDNA was isolated from pSP653 (Gerlach et al. 1985, Virology 151: 172-185) and ligated to Acc I - Spe I cut, phosphatase-treated pGEM 4 to reform the Acc I site. A resulting clone was linearized with Acc I, phosphatase-treated and a 359 bp Taq I fragment of the sTobRV cDNA was inserted. The resulting clones were screened for the presence of a circularly permuted 520 bp sTobRV cDNA sequence containing the terminally redundant residues 277 to 81 (pTTS). The sTobRV sequence is flanked by promoters for T7 and SP6 RNA polymerases, and in vitro transcription gave rise to RNAs of (+) or (-) orientation which contained two sites for self-cleavage.

In vitro Mutagenesis:

The plasmid pTTS (50 ug) was linearized with BamHI, treated with S1 nuclease and religated, to remove a unique BamHI site. The resulting construction,

pTTS-B, was treated with 2×10^{-4} units DNase I in 20mM Tris-HCl pH 7.0, 15 mM $MnCl_2$ for 10 mins. at 37°C. The resulting linear DNAs were trimmed and/or end-filled using T4 DNA Polymerase, and purified by 0.7% LGT agarose gel electrophoresis and extraction. Kinased BamH I linker sequences (CGGATCCG) were ligated to the linearized plasmid overnight at room temperature in the presence of 5% polyethylene glycol. Subsequently, the reactions were BamH I digested, and the linear plasmid DNAs repurified by 0.7% LGT agarose gel electrophoresis (this was found necessary to remove last traces of circular plasmid, together with unligated linkers). Plasmids were recircularized using T4 DNA ligase and transformed into E. coli DH-1. Colonies (greater than 1000) were scraped from agar plates, grown in liquid culture to saturation and a mixed population of plasmid DNAs prepared. The mixed sTobRV cDNA inserts were excised by restriction enzyme digestion at flanking EcoRI and PstI sites, purified by 1% LGT agarose gel electrophoresis, and sub-cloned into EcoRI - PstI cut, phosphates-treated pGEM 4. The resulting transformants were again pooled, grown in liquid culture and plasmid DNA prepared. The plasmid DNAs were treated with BamHI, to cleave only those plasmids containing a BamHI linker sequence, and the linear forms were again purified by two rounds of 0.7% LGT agarose gel electrophoresis, recircularized with T4 DNA ligase, and transformed into E. coli DH-1. Individual transformants were screened for the approximate position of the inserted BamHI linker within the sTobRV sequence by restriction enzyme digestion, sub-cloned into M13 mp19 and sequenced via the dideoxynucleotide chain termination technique.

A library of sTobRV mutants resulted, and nucleotide sequence analysis showed that each mutant contained an inserted BamHI linker sequence (CGGATCCG) together with flanking duplicated or
5 deleted sTobRV sequences. The mutants were transcribed in vitro and the RNAs assayed for their ability to undergo cleavage. From these experiments, a 52-nucleotide sequence was identified as containing both the substrate and cleavage portions of sTobRV
10 RNA. This 52-nucleotide sequence, depicted in Figure 1b, contained the domain of conserved sequence required for self-cleavage of other RNAs (Figure 1a). One mutant, designated D-51, contained an eight nucleotide BamHI linker sequence inserted between
15 three duplicated sTobRV nucleotides numbered 7 to 9. This mutant underwent self-catalysed RNA cleavage.

97 and 108 base-pair HaeIII fragments containing the 52-nucleotide cleavage sequence of the wild type and D-51 RNAs (as shown in Figures 1b and 1c) were
20 excised from sequenced plasmid clones. The fragments were ligated into the SmaI site of pGEM4 and screened to obtain both orientations of the insert. The plasmids were linearised using EcoRI and (+) and (-) strand RNAs of lengths 159 and 170 bases were
25 transcribed using 200 units/ml T7 RNA polymerase in 50 mM Tris-HCl, pH 7.5, 10 mM NaCl, 6 mM MgCl₂, 2 mM spermidine, 1000 units/ml RNasin, 500 µM ATP, CTP and GTP with 200 µM [³²P] UTP. RNAs were fractionated by electrophoresis on a 10%
30 polyacrylamide, 7 molar urea, 25% formamide gel, and autoradiographed.

As shown in Figure 1d, no cleavage of the (-) strand RNA transcripts was observed. This was as expected, as the (-) strand did not contain a

self-catalysed cleavage site. With the (+) strands of both the wild type and D-51 sequences, cleavage took place, with cleavage of the D-51 RNA being somewhat less efficient than that of the wild type (Figure 1d). This experiment indicates that the single stranded loop region at the right-hand side of the 52-nucleotide sequence involved in the self-catalysed cleavage of RNA is non essential.

Separation of Enzymic and Substrate Activities:

Using the BamHI restriction endonuclease site inserted into D-51, the flanking HaeIII-BamHI and BamHI-HaeIII fragments were obtained and each was sub-cloned into an E. coli plasmid suitable for in-vitro transcription. This led to the elimination of the mutated single-stranded loop from the self-cleavage domain, splitting the region into two RNA segments (Figure 2a). The smaller HaeIII-BamHI fragment contained nucleotides 321 to 9, including the actual site of cleavage and was termed the S fragment. The BamHI-HaeIII fragment containing sTobRV nucleotides 7 to 48 was termed the ribozyme or Rz fragment. The E. Coli plasmids used for in-vitro transcription were pGEM3 and pGEM4 (Promega, Madison, WI, U.S.A.). These expression plasmids contain:

- (a) an origin of replication;
- (b) selectable drug resistance (Amp^r) gene;
- (c) a multiple cloning site flanked by RNA polymerase promoters which can be used for in vitro production of transcripts.

T7 DNA polymerase treated, KpnI digested Rz-pGEM3 and XbaI digested S-pGEM4 were transcribed using SP6 RNA polymerase under the same conditions as set out above.

As shown in Figure 2, both the S and Rz-RNAs showed no significant degradation when incubated alone (Figures 2b, lanes 1 and 3) under conditions suitable for highly efficient self-cleavage (50°C, 20mM MgCl₂, pH 8.0). The labelled Rz-RNA also appeared unaltered after incubation with the S-RNA (Figure 2b, lanes 2 and 5). However, when the S-RNA was mixed with the Rz-RNA, efficient cleavage of the S-RNA occurred (Figure 2b, lanes 4 and 5) producing two fragments. The product sizes were consistent with cleavage of the S-RNA (84 bases) at the normal site between nucleotides #359 and #1, to give 5' and 3' proximal fragments of 67 and 17 nucleotides, respectively. This shows that the S-RNA acted as a substrate for ribonucleolytic cleavage by the Rz-RNA, which acted in a catalytic fashion.

A model of a ribozyme based on the catalytic region of sTobRV RNA is shown in Figure 3. The ribozyme has two arms or flanking sequences of single stranded RNA shown at C, hybridizing to complementary sequences on a substrate RNA, i.e. RNA to be cleaved. Each flanking sequence shown at C, contains 8 ribonucleotides. The number of nucleotides contained in region C is not critical. Sufficient nucleotides must, however, be present to allow the ribozyme to hybridize to a target RNA. Four nucleotides in each region C appears to be the minimum number for hybridization.

The catalytic region B contains sequences which are highly conserved in naturally-occurring cleavage domains (see Figure 1a). From a comparison with cleavage domains of the known sequences, the length of the base pair stem II is unimportant, as is the presence of an associated loop at one end thereof.

The cleavage site within the target RNA is depicted at A (in Figure 3) as GUC. On the basis of our experiments (not shown), and others by Koizumi (FEBS LETT 288; 228-230 (1988); and FEBS LETT 239; 285-288 (1988)) on the cleavage sites in naturally occurring RNAs, the sequences GUA, GUC, CUC, AUC and UUC also act as cleavage sites within RNA.

EXAMPLE 2

10 Demonstration of the Design, Synthesis and Activity of Ribozymes with New and Highly Specific Endoribonuclease Activity:

As an illustration of this invention, three ribozymes have been designed, which are targeted against the transcript of a commonly used indicator gene derived from bacteria, Tn9 Chloramphenicol Acetyl Transferase (CAT), which can provide antibiotic resistance in bacteria, plants and animals and can be easily assayed. These ribozymes, designated RzCAT-1 to 3 correspond to potential GUC cleavage sites in CAT RNA at positions 139-140, 494-495 and 662-663 respectively. The sequences of these ribozymes are depicted in Figure 4. In each case, the flanking sequences of the ribozyme which hybridize to the target CAT RNA, were 8 nucleotides in length. The catalytic region was chosen to correspond to that of sTobV RNA, shown in Figure 3.

The CAT gene was obtained from pCM4 and sub-cloned as a BamHI fragment into pGEM-32 (from Promega, Madison, WI, U.S.A.). This plasmid was linearised with HindIII and CAT gene transcripts were obtained using T7 RNA polymerase with 220uM [α -³²P]UTP. Ribozyme sequences were synthesised

as oligodeoxynucleotides, Rz CAT-1, 2 and 3, respectively. They were kinased, ligated with phosphatased treated, EcoRI-PstI cut pGEM4 and incubated with the Klenow fragment of DNA polymerase 5 1 before bacterial transformation. EcoRI linearised plasmids were transcribed with T7 RNA polymerase to produce ribozyme RNAs. Ribozymes were incubated with CAT transcript in 50 mM Tris-HCl pH 8.0, 20 mM $MgCl_2$ at 50°C for 60 min, and the products 10 fractionated by 5% polyacrylamide 7M urea, 25% formamide gel electrophoresis prior to autoradiography.

When the 840-nucleotide CAT transcript was incubated with any one of the three ribozymes, 15 efficient and highly sequence-specific cleavage occurred (Figure 5) producing 2 RNA fragments in each reaction. The fragment sizes were consistent with the predicted sites for cleavage (i.e. 139 and 696, 494 and 341, 662 and 173 base fragments were the 5' 20 and 3' products from RzCAT-1 to 3 catalysed cleavage respectively). The conditions required for these ribozyme-catalysed cleavages were similar to those observed for naturally occurring cleavage reactions (Foster, A.C. and Symons, R.H., Cell 49: 211-220 25 (1987) and Foster, A.C. and Symons, R.H. Cell 50: 9-16 (1987)), with more efficient cleavage occurring at elevated pH, temperature and divalent cation concentrations (data not shown). When present in molar excess, the three ribozymes catalysed almost 30 complete cleavage of the CAT RNA substrate after 60 min. in 50 mM Tris HCl, pH 8.0, 20 mM $MgCl_2$ at 50°C. Under similar conditions with 0.1 μM substrate and 3 μM ribozymes, the $T_{1/2}$ of CAT mRNA substrate was 3.5, 3.5 and 2.5 min. in the

presence of RzCAT-1 to 3 respectively. The ribozyme sequences were inactive against the complement of the substrate RNA (i.e. the (+) strand), and in the form of oligodeoxyribonucleotides (data not shown). The
5 3' terminal cleavage fragments from each ribozyme catalysed reaction were isolated and 5' ^{32}P -kinased (50 mM TrisHCl pH 9, 10 mM MgCl_2 , 10 mM DTT with 50 uCi γ - ^{32}P ATP and 5 units T4 polynucleotide kinase for 30 min. at 37°C.).
10 Efficient kinasing of the fragments indicated that they possessed 5' terminal hydroxy groups, similar to those produced in naturally occurring cleavage reactions.

The terminal nucleotide of the fragments produced
15 by cleavage of the CAT sequences by RzCAT-1 to 3 were determined. Briefly, radiolabelled fragments were purified on a 5% polyacrylamide gel and digested with an equal volume of 500 units/ml RNase T1, 25 units/ml RNase T2 and 0.125 mg/ml RNaseA in 50 mM ammonium
20 acetate pH 4.5 for 120 min. at 37°C. The products were fractionated on a 20% polyacrylamide gel containing 25 mM sodium citrate, pH 3.5 and 7 molar urea. Figure 5b shows that the cleavage of the CAT sequences by RzCAT-1 to 3 occurs precisely before
25 nucleotides A, U and G respectively.

The terminal sequence of the CAT gene fragments were determined directly using the partial enzymatic digestion technique (Donis-Keller et al., Nucleic Acids Res. 4: 2527-2538 (1980)), using base-specific
30 partial ribonucleolytic cleavage. The sequence of the fragments confirmed that cleavage occurred at the expected locations within CAT RNA (not shown).

Enzymatic Catalysis:

To demonstrate that ribozymes cause cleavage of

the CAT mRNA substrate in a catalytic manner, each was incubated with a molar excess of substrate, under conditions which should favour both efficient cleavage and product dissociation. Figure 6 shows the results of an experiment where after 75 min. at 50°C, pH 8.0 in 20 mM MgCl₂, 10 pmoles of RzCAT-1 had catalysed specific cleavage of 163 pmoles of a truncated CAT mRNA (173 bases) substrate to give 5' and 3' fragments of 139 and 34 bases, respectively. On average, each ribozyme had participated in greater than ten cleavage events. After 75 minutes at 50°C some non-specific cleavage of RNA was noticed due to the extreme conditions, but 70% of the remaining intact RNAs (163 pmoles) had accumulated as the 139 base fragment. Similar results were obtained for RzCAT-2 and 3 (data not shown), and thus each acts as an RNA enzyme.

EXAMPLE 3

20 The Effect of Temperature on Ribozyme Activity:

The effect of reaction temperature on the in-vitro rate of ribozyme activity was examined.

A time course of reactions for ribozymes RzCAT-1 to 3 on CAT RNA substrate at 37°C and 50°C was carried out.

In this experiment, reactions for each ribozyme were set up in duplicate, using reaction conditions for ribozyme cleavage set out in Example 2. One reaction was incubated at 37°C, the other at 50°C. Samples were removed at time points to 90 minutes and the extent of reaction was analysed by denaturing polyacrylamide gel electrophoresis. Figure 7 shows the time course of reaction for each of the ribozymes RzCAT-1 to 3 at 37°C and 50°C. The reaction rate of

each ribozyme increases with increased reaction temperature.

The time taken for 50% ($t_{1/2}$) cleavage of CAT RNA is set out in Table 1.

5

TABLE 1

	RzCAT-1	RzCAT-2	RzCAT-3 $t_{1/2}$ (mins.)
50°C	3.5	3.5	2.5
10 37°C	55.0	70.0	65.0

As shown in Table 1, the rate of reaction of ribozymes at 37°C is approximately 20 times slower than the rate of reaction at 50°C.

15

EXAMPLE 4

The Effects of Varying Arm Lengths of Ribozymes (or Flanking Sequence) on Ribozyme Catalytic Activity:

20 The arms or flanking sequences of a ribozyme (region (I) of formula 1) hybridize the ribozyme to a target RNA whereafter cleavage of the RNA takes place. In this experiment, the effect on cleavage rate of a target sequence by altering the extent of complementarity and subsequent length of base pairing
25 of the ribozyme arms to the target sequence was investigated.

Ribozymes were produced with 4, 8 and 12 base complementarity to the target sequence RzCAT-2 on each arm (Figure 8a). The ribozymes were prepared
30 according to the methods of Example 2. Ribozyme activity was determined by incubating the ribozyme RNA with CAT RNA as described previously.

The ribozyme having a 4 base complementarity on

each arm did not cleave the substrate RNA. The ribozyme with 8 base complementarity on each arm cleaved the CAT substrate as did the ribozyme having 12 base complementarity. Ribozymes with 12 base complementarity cleaved target RNA more efficiently, as judged by reaction rate in-vitro than did ribozymes having a lesser number of base complementarities. Even though it appears necessary to have more than four base complementarity, increasing the length of the hybridizing region of ribozymes increases their reaction rate.

In a second experiment, the reaction efficiency of a ribozyme having (a) complementarity to the entire length of the CAT transcript target RNA and (b) multiple catalytic domains, was investigated.

Four GUC target sites in CAT RNA sequences were chosen. Ribozyme catalytic domains against these sites were "inserted" into a complete anti-sense (-) sequence for the CAT transcript and catalytic activity tested.

The four sites chosen were the three specified by RzCAT-1 to 3 described previously, and a further site which may be represented as follows:

New CAT site

#192

	His	His	Ala	Val	Cys	Asp	Gly	
5'	CAU	CAU	GCC	GUC	UGU	GAU	GGC	3'

where "192" refers to amino acid 192 in the CAT polypeptide and an refers to the site of cleavage.

Oligodeoxyribonucleotides containing ribozyme catalytic domains and spanning each of these cleavage sites were used for M13 mutagenesis experiments to produce a sequence containing the entire complement of the CAT sequence but with the four ribozyme

catalytic domains inserted within it. M13 mutagenesis was performed by binding of oligonucleotides containing ribozyme insertions to single stranded M13 DNAs containing uracil, followed
5 by synthesis of complementary DNAs containing the insertion. The complementary DNAs were recovered following cloning in an appropriate Escherichia coli strain, (T.A. Kunkel 1985, Proc. Natl. Acad. Sci. U.S.A. 82: 488-492). The resultant double-stranded
10 cDNA was cloned in an in-vitro expression vector to produce ribozyme RNA using the T7 polymerase transcription system. Ribozyme activity was determined by incubation of ribozyme RNA with CAT transcript followed by gel electrophoresis of the
15 reaction mixture, after glyoxal treatment to denature nucleic acids.

Autolytic cleavage occurred at all the expected sites on the CAT transcript. Accordingly, the flanking sequences of arms or a ribozyme may extend
20 along the full length of the RNA transcript which is to be cleaved.

Figure 9 schematically shows the production of catalytic anti-sense RNA containing each of the four ribozymes. The catalytic anti-sense RNA contains
25 approximately 900 bases.

Under the above reaction conditions, the ribozyme and target sequences form high molecular weight complexes presumably by extensive base pairing. A strong denaturing treatment such as glyoxal treatment
30 is required to resolve reaction products during electrophoresis.

EXAMPLE 5**Target Sequences for Ribozyme Cleavage:**

The GUA motif in mRNA was tested to see whether a ribozyme would effect RNA cleavage at this sequence.

5 A specific site in CAT mRNA, including the GUA motif (Figure 10a) was chosen and an appropriate ribozyme sequence was prepared and tested for activity. The ribozyme contained arms of 8 ribonucleotides.

10 Synthetic oligonucleotides corresponding to the ribozyme of Figure 10 were prepared according to Example 2 and double-stranded cDNA cloned into an in-vitro expression vector (pGEM 4, see above) in E. coli in order to produce ribozyme RNA using the T7
15 polymerase transcription system. Ribozyme activity was determined by incubation of ribozyme RNA with CAT mRNA, followed by gel electrophoresis of the reaction mixture as previously described.

The ribozyme effected cleavage at the GUA target
20 site (not shown). Accordingly, the motif GUA in RNA is a substrate for ribozymes of the present invention. This was not completely unexpected since one naturally occurring cleavage site in the satellite RNA of lucerne transient streak virus requires
25 recognition of a GUA site.

Similarly, a GUU motif in the CAT RNA target sequence was tested with an appropriate ribozyme (See Figure 10b), and cleavage was effected.

30 EXAMPLE 6**Ribozyme Cleavage of Viral RNA:**

Viroid RNA, in the form of citrus exocortis viroid RNA, was cleaved using a ribozyme of the present invention.

Two GUC target sites were chosen in citrus exocortis viroid (CEV) RNA. One site in the complementary strand sequence was also chosen. Ribozymes were prepared against all of these sites
5 and tested for activity. The ribozymes were designated CEV9x(+), CEV9x(-) and CEV25x(+). Figure 11 shows the three cleavage sites in CEV RNA for each of these ribozymes.

Ribozymes were prepared according to previous
10 methods. Ribozyme RzCEV25x(+) is shown in Figure 12. This ribozyme cleaves the GUC motif at nucleotide 116 of CEV RNA.

Figure 12(b) shows cleavage of CEV RNA with ribozyme RzCEV9x(+). No cleavage is observed with
15 ribozyme RzCEV9x(-).

This experiment indicates that ribozymes are active against target RNA sequences from diverse sources. This is to be expected, as all RNAs are formed from the basic ribonucleotide building blocks
20 of adenine, guanine, cytosine and uracil, regardless of their animal, plant or microbial origin.

EXAMPLE 7

Examples of Ribozymes Having Variable Catalytic Domains

25 A ribozyme targeted against a CAT-2 site was prepared using the catalytic domain sequence from the satellite RNA of subterranean clover mottle virus (SCMoV). Twelve base complementarity of ribozyme arm flanking sequence was incorporated into the design of
30 the ribozyme RzSCMoV. The ribozymes RzCAT-2 and RzSCMoV are shown at Figures 13a and 13b respectively. The loop region of RzSCMoV contains 5 nucleotides, having the sequence AAAUC. This is to be contrasted with the loop region of RzCAT-2 which

contains 4 nucleotides having the sequence AGAG. In addition, RzSCMoV contains a C in the catalytic region, in place of U* in RzCAT-2. The different sequences in RzSCMoV when compared with RzCAT-2 are 5 marked.

The RzSCMoV was produced according to Example 2. RzSCMoV was active, yielding two cleavage products as expected.

In another experiment, the citrus exocortis 10 viroid (CEV) target site at nucleotide -336 in its complementary RNA was cleaved using a ribozyme (RzCEV2) having the sequence set out in Figure 14a. The loop region designated with the letter "L" in Figure 14 comprises six nucleotides having the 15 sequence 3'-CCTATA-5'. This is distinct from the loop region of sTobrV which comprises four nucleotides having the sequence 3'-AGAG-5'. This ribozyme cleaves target CEV complementary RNA at position -336 as shown in the electrophoretic profile 20 of Figure 14b.

This experiment indicates that the number of nucleotides, and nucleotide sequence of the loop region is unimportant in ribozyme activity. In these experiments, the ribozyme was produced according to 25 methods previously described in the specification.

In another experiment, the effect of base pairing in the catalytic domain (stem region) on ribozyme activity was investigated.

A modified ribozyme corresponding to RzCAT-2 but 30 containing four extra base pairs was prepared and tested. In Figure 15a, the sequence of the ribozyme Rz CAT-2 is shown hybridized to target CAT RNA. The test ribozyme is shown at 15b, with the additional base pairs boxed. The test ribozyme had comparable

activity to that of RzCAT-2. This indicates that the base paired region of the ribozyme catalytic domain may be of variant length, without effecting catalytic activity.

- 5 We have observed (data not shown) that the stable in vivo form of sTobRV RNA transcripts expressed in transgenic plants is primarily circular, presumably due to ligation of 5' and 3' termini. Therefore, the use of two autolytic cleavage sites flanking a
10 sequence of interest in an in vivo RNA transcript is likely to lead to a circularized product which may have greater stability than linear transcripts. This approach appears to provide a novel method for in vivo stabilization of ribozyme sequences. This is
15 termed circularization.

EXAMPLE 8

In-vivo Activity of Ribozymes:

- The in-vivo activity of ribozymes in plant cells
20 is investigated in this Example.

Experimental Protocol:

- Plasmids containing anti-CAT (CAT=chloramphenicol acetyl transferase) or combined anti-CAT/ribozyme gene constructions (see below) were introduced into
25 tobacco protoplasts in the same amount and proportion relative to each other, along with another plasmid which contained a functional CAT gene construction. CAT activities were measured and compared with the base level of gene activity.

30 Materials and Methods:

- (a) Electroporations and CAT assays.


These were performed as described in Llewellyn et al. J. Mol. Biol. (1987) 195:115-123. Briefly, protoplasts of Nicotiana plumbaginifolia line T5 were

prepared from a suspension two days after subculture, suspended in 10mM HEPES, pH 7.2, 150mM NaCl, 0.2M mannitol and adjusted to a density of 3×10^6 /ml. Electroporation was carried out using a single 50ms pulse at 250V. Protoplasts were diluted 10-fold and cultured for 20 hr. at 26°C in the dark. They were disrupted by sonication and extracts obtained. The extracts were normalized for protein content and assayed for CAT activity *in vitro* using ¹⁴C-chloramphenicol and acetyl CoA. Reaction products were separated by thin layer chromatography and visualized by autoradiography. Extent of reactions were calculated by the production of radioactive product derivatives from the ¹⁴C-chloramphenicol template.

(b) Gene Constructions.

Gene constructions were introduced into 0.1 ml protoplast suspensions as plasmid DNAs which had been purified from bacteria by extraction and two cycles of CsCl equilibrium density gradient centrifugation. They were resuspended in 10 mM Tris/1 mM EDTA/pH=7.5 for use.


The active CAT gene construction was borne on the plasmid designated pCAT7+. It was derived by fusion of a CAT gene sequence (from plasmid pCM4, see T.J. Close and R. Rodriguez, 1982, Gene 20:305-316) into the plasmid pJ35SN (derived from p35SN, W.L. Gerlach et al., 1987, Nature 328:802-805) so that the active gene construction was:




35S refers to the 35S CaMV (cauliflower mosaic virus) promoter, NOS to nopaline synthetase polyadenylation signal, T/C to transcription.

Along with 0.2 ug of pCAT7+ there were added various gene constructions in excess as described below. The gene constructions were contained within plasmids with the following designations:

- 5 pJ35SN - This vector plasmid, a map of which is shown in Figure 16, contains a 35S CaMV promoter and plant nopaline synthase 3' polyadenylation signal, which may be depicted as follows:



- pCAT7- - This contains the CAT gene sequence inserted into pJ35SN such that transcription will result in the production of the antisense CAT RNA, which may be depicted as follows:



- pCAT19- - This contains the CAT gene with four catalytic ribozyme domains included within it, (see Example 4 and Figure 9), inserted into pJ35SN such that transcription will result in the production of antisense CAT RNA containing four catalytic ribozyme domains, which may be depicted as follows:



catalytic domain insertions

Results:

- 30 The following table shows the relative CAT activities in cells 20 hours after electroporation. Activity is expressed as per cent conversion of chloramphenicol substrate in a 1 hour assay.

		ug Plasmid Electroporated				% Conversion
Treatment		pCAT7+	pJ35SN	pCAT7-	pCAT19-	
5	1A					0
	1B					0
	2A	0.2	18			21
	2B	0.2	18			46
	3A	0.2	9	9		28
	3B	0.2	9	9		32
	4A	0.2		18		26
	4B	0.2		18		19
	5A	0.2	9		9	19
	5B	0.2			9	22
10	6A	0.2			18	14
	6B	0.2			18	16

15 (for each treatment "A" and "B" are duplicates)

The following conclusion can be drawn from these results:

- 20 (a) The introduction of the CAT gene construction results in significant CAT activity - compare 2A,B with 1A,B. There is variability between duplicates. From the trends seen in the other samples (see "b" and "c" below) it is likely that 2A shows an abnormally low activity.
- 25 (b) Concomitant introduction of an antisense gene construction results in a decrease in the level of activity - compare 3A,B and 4A,B with 2B. The extent of the decrease is related directly to the level of the antisense gene added as plasmid - compare 3A,B and 4A,B.
- 30 (c) Concomitant introduction of the combined antisense/ribozyme gene construction results in a decrease in gene activity - compare 5A,B and 6A,B with 2B. Furthermore, the decrease is more

marked than for the corresponding levels of antisense gene constructions - compare 5A,B with 3A,B, and 6A,B with 4A,B.

The average results for four in-vivo experiments are shown in Figure 17. In this Figure, "control" represents treatment 2. "Antisense" represents treatment 4. "Catalytic" represents treatment 6 and "Background" represents treatment 1.

The catalytic ribozyme inhibits CAT activity an average of 47%, compared to an average of 34% for an antisense ribozyme.

The introduction of ribozyme-bearing genes into plant cells inhibits the activity of genes against which they are targeted. Furthermore, the inhibition is greater than for corresponding antisense RNA molecules.

These results show that ribozymes will be active in animal, plant or microbial cells against a range of target RNA molecules.

The mechanisms of action of the ribozymes in this Example is unclear. For example, the antisense ribozyme may irreversibly hybridize to a target RNA and catalyse phosphodiester bond cleave at one or more selected target sites along the target RNA. Alternatively, cellular enzymes may unwind the antisense RNA from its target sequence, such that the target RNA is cleaved into two or more fragments.

EXAMPLE 9

In-vivo Activity of Ribozymes in Animal Cells:

The activity of ribozymes in inactivating a target RNA in mammalian cells is demonstrated in this

5 Example.

Materials and Methods:

The active gene constructions encoding ribozymes were transfected into the widely available monkey kidney cell line COS1 by electroporation. In this
10 method, 3×10^6 /ml COS1 cells suspended in a buffered saline with 10% FCS (foetal calf serum), were contacted with various gene constructs and an electric discharge applied to effect electroporation of DNA into the cells. The transfected cells were
15 incubated at 37°C for 48 hours in culture medium before assay for CAT and luciferase activity.

CAT gene constructs were borne on the plasmid designated pTK CAT (Miksicek et al., Cell 46: 283-290, 1986). This plasmid was derived by the
20 introduction of a CAT gene sequence into the plasmid pSV2 such that it is under the control of the thymidine kinase promoter of the herpes simplex virus.

Gene constructs encoding ribozymes were borne on the plasmid pSV232A (De Wet et al., Molecular and
25 Cellular Biology 7: 725-737, 1987) containing the luciferase gene fused to the SV40 early promoter. DNA encoding ribozymes was ligated into the XbaI site at the 3' end of the luciferase gene according to the standard methods of Maniatis et al. (Molecular
30 Cloning, A Laboratory Manual, Cold Spring Harbour, 1982).

The following constructs were prepared using standard techniques of Maniatis et al. (Supra):
pFCS8 - This plasmid vector contains DNA encoding

ribozyme RzCAT-1 fused to the 3' end of the luciferase gene in a non-functional orientation.

This may be depicted as follows:



where 232A refers to pSV232A sequences, SV40 early refers to the early promoter of SV40 and small T is DNA encoding the small T intervening sequence of SV40. This construct results in production of an RNA molecule encoding luciferase and the ribozyme RzCAT-1, the latter being in an orientation such that it would not be expected to be catalytic.


15 pFC4 - This plasmid is the same as pFC58 except that RzCAT-1 is replaced with RzCAT-3.

pFC1-6 - This plasmid is the same as pFC58 except that RzCAT-1 is replaced with RzCAT-3 in the sense orientation (5'-3').

20 pFC20 - This plasmid is the same as pFC1-6 except that RzCAT-3 is replaced with RzCAT-2 having eight nucleotide flanking sequences.

pFC12 - This plasmid is the same as pFC20 except that the ribozyme RzCAT-2 contains twelve nucleotide flanking sequences.

25 pFC50 - This plasmid contains the CAT gene with four catalytic ribozyme domains included within it (see Example 4 and Figure 9) in the sense orientation (5'-3'), which on transcription gives rise to an inactive ribozyme. This plasmid may be depicted as follows:



- pFC54 - This plasmid is the same as pFC50, except that the CAT gene and the ribozyme domains are in the antisense (3'-5') active orientation.
- 5 pFC64 - This plasmid shares the SV40 promoter and polyadenylation signals with pFC50 and contains the wild type CAT gene with no inserted ribozyme domains. This gene is in an antisense orientation, and thus does not
- 10 produce CAT protein.
- pFC65 - This plasmid is the same as pFC64 except that the wild type CAT gene is in the sense (5'-3') orientation and is thus productive of CAT protein.

15 Assays:

Luciferase activity was assayed according to the methods of De Wet et al. (Supra). Briefly, COS cells were lysed 48 hours after transfection, and the cell lysate incubated with luciferin, the substrate of

20 luciferase, and luminescence detected using a scintillation counter.

CAT activity was also measured using COS cell lysates (cell lysates were divided into two, and each portion assayed either for luciferase or CAT

25 activity), according to the method of Sleight, M.J., Anal. Biochem. 156: 251-256, (1986).

In the in-vivo assays, pFC58 and pFC4 did not effect CAT activity in transfected cells. This activity was designated 100% CAT activity and 0% CAT

30 suppression. CAT activity in cells transfected with other plasmids was measured relative to pFC58. The % of CAT suppression was measured as

$$100 - \frac{39 \left(\frac{\text{CAT}_{\text{test}}}{\text{CAT}_{\text{control}}} \right) \times 100}{\text{CAT}_{\text{control}}}$$

normalised to luciferase production. CAT_{test} = CAT assay result for test constructs. CAT control = CAT assay for control constructs (pFC4 and pFC58).

Luciferase production is an internal control for electroporation, and gives a measure of ribozyme production within each individually electroporated tissue culture plate.

Results:

Experiment (i)

µg plasmid electroporated/ 1.5×10^6 cells						
Treatment	PTKCAT	pFC58	pFC20	pFC1-6	pFC12	% CAT Suppression
1	5		2			56
2	5		1	1		53
3	5				2	40
4	5	2				0

All treatments were carried out in duplicate and an average value given.

Experiment (ii)

µg plasmid electroporated/ 1.5×10^6 cells						
Treatment	PTKCAT	pFC-1-6	pFC20	pFC12	pFC4	% CAT Suppression
5	5	2				75
6	5		2			75
7	5		4			62
8	5	1	1			70
9	5			4		51
10	5				2	0

Treatments 5 to 10 were carried out in duplicate and an average value given.

Experiment (iii)

µg plasmid electroporated/ 1.5×10^6 cells				
Treatment	PTKCAT	pFC1-6	pFC4	% CAT Suppression
11	5	2		66
12	5		2	0

Treatments were carried out in quintuplicate and an average value given.

Experiment (iv)

Treatment	pTKCAT	µg plasmid electrophorated				% CAT Suppression
		pFC50	pFC54	pFC64	pFC65	
13	5	2				0
14	5		2			26
15	5			2		2
16	0				2	NA

Each of treatments 13 to 16 were carried out in quadruplicate.

The sense CAT construction (treatment 16) was productive of high levels of CAT activity in its own right. Therefore % suppression is not applicable (NA).

A number of experiments were also conducted with the TK promoter of pTKCAT replaced with the human metallothionein promoter. When this CAT construct was co-transfected into COS1 cells with plasmids encoding one or more ribozymes, a marked decrease in CAT activity was observed.

The above results clearly demonstrate the in-vivo inactivating activity of ribozymes in animal cells.

Whilst the effectiveness of the ribozymes in vivo, is believed to be caused by one or more catalytic regions which are capable of cleavage of a target RNA, the presence of such regions in "Antisense" RNA type ribozymes may not actually lead to cleavage in vivo if the entire RNA/antisense RNA molecule does not fall apart. However, regardless of whether the molecule falls apart or not, the foregoing Examples demonstrate the effectiveness of the ribozyme in inactivating the target RNA. Thus, the invention is applicable to all ribozymes having a catalytic region capable of causing cleavage and a hybridizing region, regardless as to whether cleavage

41

actually occurs in the target RNA. i.e. The hybridizing region may be so large as to cause the combined RNA/ribozyme to stay together and prevent the target RNA being cleaved into separated
5 components even though the catalytic region is itself capable of causing cleavage.

10

CLAIMS:


1. A ribozyme comprising a hybridizing region which is complementary in nucleotide sequence to at least part of a target RNA, and a catalytic region which is adapted to cleave said target RNA wherein the hybridizing region contains 9 or more nucleotides.

2. A ribozyme comprising one or more arms formed of single stranded RNA and having a sequence complementary to at least part of a target RNA, said one or more arms being associated with a catalytic region, capable of cleaving said target RNA; and where the hybridizing region comprises a single arm of RNA said arm contains at least 9 nucleotides, and where the hybridizing region comprises 2 or more arms of RNA, the sum of nucleotides in said arms is greater than 9 nucleotides.

3. A ribozyme according to claim 2, having two single stranded arms of RNA, associated with one another through said catalytic region.

4. A ribozyme according to any preceding claim which does not modify cleaved target RNA.


5. A ribozyme of the formula:



wherein; X represents any ribonucleotide and each X residue may be the same or different;
the sum of n and n' is greater than 6 and n and n' may be the same or different;
an (*) represents a base pair between complementary ribonucleotides;
X' and X'' represent ribonucleotides of complementary sequence along at least part of their length to allow base pairing between the oligoribonucleotides, or X' and X'' together form a single RNA sequence wherein at least part of said sequence comprises a stem formed by base pairing between complementary nucleotides;
and optionally, an additional nucleotide selected from any one of A, G, C or U is inserted after ¹A.

6. A ribozyme as claimed in any preceding claim having the formula:

44



wherein; X represents any ribonucleotide and each X residue may be the same or different;
 an (*) represents a base pair between complementary ribonucleotides;
 n and n' are as previously defined;
 B represents a bond, a base pair, a ribonucleotide, or an oligoribonucleotide containing at least 2 ribonucleotides;
 m and m' are 1 or more and may be the same or different; and
 optionally, an additional nucleotide selected from any one of A, G, C or U is inserted after ¹A.

7. A ribozyme according to any one of claims 5 or 6, where the sum of n and n' is 16 or greater.

8. A ribozyme according to any preceding claim wherein the catalytic region is capable of cleaving a target RNA after the sequence X⁰UY where X⁰ is any ribonucleotide, U is uracil and Y is adenine, cytosine or uracil.

9. A ribozyme containing multiple catalytic domains connected through flanking sequences, each said flanking sequence including sequences complementary to the target RNA which is desired to be inactivated.

10. A ribozyme according to any preceding claim which has been stabilized by circularization.

11. A ribozyme comprised of catalytic antisense sequences according to any preceding claim.

12. A method for the inactivation or cleavage of a target RNA which comprises reacting said target RNA with a ribozyme as claimed in any one of claims 1 to 11.

13. A method according to claim 12, wherein said inactivation occurs in-vivo in prokaryotic or eukaryotic cells, as a result of transcription of said ribozyme from a DNA or RNA transfer vector transfected into said cells or precursors thereof, or expression of a gene encoding said ribozyme which is integrated into the host genome, such that said ribozyme hybridizes to and inactivates a desired cellular target RNA.

14. A method according to claim 13, wherein said cells are plant or animal cells.

15. A method according to claim 13 or 14, wherein the target RNA is endogenous to the cells, or is exogenous, such as viral RNA.

16. A method according to claim 15, wherein target viral RNA is selected from RNA of the HIV(AIDS), hepatitis or other virus causing disease in mammals.

17. A method according to claim 15, wherein the viral RNA is bacteriophage RNA.

18. A method for the treatment of viral disease in man or animals which comprises administering to a subject in need of such treatment, an effective amount of a ribozyme according to any one of claims 1 to 11, said ribozyme hybridizing to and inactivating transcripts of said viral RNA.

19. A method for treating viral or viroid disease in plants which comprises administering to a plant a ribozyme according to any one of claims 1 to 11, said ribozyme hybridizing to and inactivating transcripts of said viral RNA.

20. A composition which comprises a ribozyme as claimed in any one of claims 1 to 11 in association with pharmaceutically, veterinarily, or agriculturally acceptable carrier or excipient..

21. A method for producing a ribozyme according to any one of claims 1 to 11 which comprises the steps of:

- (a) ligating into a transfer vector comprised of DNA, RNA or a combination thereof a nucleotide sequence corresponding to said ribozyme;
- (b) transcribing the nucleotide sequence of step (a) with RNA polymerase; and



(c) optionally purifying the ribozyme.

22. A transfer vector comprised of RNA or DNA or a combination thereof containing a nucleotide sequence which on transcription gives rise to a ribozyme as claimed in any one of claims 1 to 11.

23. A transfer vector according to claim 22, which is a bacterial plasmid or phage DNA.

24. A prokaryotic or eukaryotic cell containing nucleotide sequences which on transcription thereof give rise to a ribozyme as claimed in any one of claims 1 to 11.

2/19



3/19




FIG. 3.

4/19




Fig. 4a.

RzCAT-1 SITE:

#24




Fig. 4b.

5/19

RzCAT-2 SITE:




Fig. 4c.

RzCAT-3 SITE:





Fig. 4d.

6/19

*Fig. 5a.*

7/19




Fig.5b.

9/19






FIG. 7.

10/19






Fig.7. (cont.) —●— SUBSTRATE —+— 5' PRODUCT

11/19





FIG. 8d.

12/19




Fig. 8a. (cont.)

13/19




FIG. 9.

14/19




Fig. 10a.




Fig. 10b.

16/19




Fig. 13a.




Fig. 13b.

17/19




Fig. 14a.




Fig. 14b.

18/19




Fig.15a.




Fig.15b.




FIG. 16.




FIG. 17
SUBSTITUTE SHEET

REVISED
VERSION

INTERNATIONAL SEARCH REPORT (SUPPLEMENTARY)

International Application No. PCT/AU 88/00478

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) 6

According to International Patent Classification (IPC) or to both National Classification and IPC Int. Cl."

II. FIELDS SEARCHED

Minimum Documentation Searched 7

Classification System | Classification Symbols

Documentation Searched other than Minimum Documentation
to the extent that such Documents are Included in the Fields Searched 8

III. DOCUMENTS CONSIDERED TO BE RELEVANT 9

Category*	Citation of Document, with indication, where appropriate, of the relevant passages 12	Relevant to Claim No 13
-----------	------------------------------------------------------------------------------------------	----------------------------

X	A.C. Forster and R.H. Symons Cell, volume 50 pp.9-16 3 July 1987 (03.07.87) "Self Clearance of Virusoid RNA is performed by the proposed 55 nucleotide active site "(See p9 and figures 4-8)	(1-3)
P Y	WO,A, 88/4330 (University Patents Inc.,) 16 June 1988 (16.06.88)	(1-4)
Y	Beer M.D. and Cech T.R. Cell, volume 47 pp.207-216, 24 October 1986 (24.10.86), "One binding site determines sequence specificity of Tetrahymena Pre-rRNA self-splicing, trans- splicing and RNA enzyme activity. (figure 1)	(1-4)

CONTINUED

* Special categories of cited documents: 10	*T* Later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step
"E" earlier document but published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Z" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

IV. CERTIFICATION

Date of the Actual Completion of the
International Search

Date of Mailing of this International
Search Report

12 DECEMBER 1989

International Searching Authority

Signature of Authorized Officer

Australian Patent Office

J.P. CHAN

FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

Y	Waring R.B. et al. Cell volume 40 pp.371-380 February 1985 (02.85) "The tetrahymena tRNA intron self-splices in E.Coli: In vivo evidence for the importance of key base-paired of RNA for RNA enzyme function". (See figures 3 and 5)	(1-4)
---	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-------

V. ☐ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE 1

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claim numbers ..., because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claim numbers ..., because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claim numbers ..., because they are dependent claims and are not drafted in accordance with the second and third sentences of PCT Rule 6.4 (a):

VI. ☐ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING 2

This International Searching Authority found multiple inventions in this international application as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.
2. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:
3. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:
4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the International Searching Authority did not invite payment of any additional fee.

Remark on Protest

- ☐ The additional search fees were accompanied by applicant's protest.
☐ No protest accompanied the payment of additional search fees.

ANNEX TO THE INTERNATIONAL SEARCH REPORT ON
INTERNATIONAL APPLICATION NO. PCT/AU 88/00478

This Annex lists the known "A" publication level patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent Document
Cited in Search
Report

Patent Family Members

WO 88/04300

EP 291533

END OF ANNEX

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁴ : C12N 9/22, 5/02, 15/00 C12N 1/20, C12P 19/34 C07H 21/02, A61K 31/73, 39/235 A61K 39/285</p>	<p>A1</p>	<p>(11) International Publication Number: WO 89/ 05852 (43) International Publication Date: 29 June 1989 (29.06.89)</p>
<p>(21) International Application Number: PCT/AU88/00478 (22) International Filing Date: 14 December 1988 (14.12.88) (31) Priority Application Numbers: PI 5911 PI 9950 PJ 0353 PJ 1304 PJ 1333 (32) Priority Dates: 15 December 1987 (15.12.87) 19 August 1988 (19.08.88) 9 September 1988 (09.09.88) 4 November 1988 (04.11.88) 7 November 1988 (07.11.88) (33) Priority Country: AU (71) Applicant (for all designated States except US): COMMON-WEALTH SCIENTIFIC AND INDUSTRIAL RE-SEARCH ORGANISATION [AU/AU]; Limestone Avenue, Campbell, ACT 2601 (AU). (72) Inventors; and (75) Inventors/Applicants (for US only) : HASELOFF, James, Phillip [AU/AU]; 17 Hemmant Street, O'Connor, ACT 2601 (AU). GERLACH, Wayne, Lyle [AU/AU]; 22 Goble Street, Hughes, ACT 2605 (AU). JENNINGS, Philip, Anthony [GB/AU];</p>		<p>13 Lobelia Street, West Chatswood, NSW 2066 (AU). CAM-ERON, Fiona, Helen [AU/AU]; 2 Rounce Avenue, Forest-ville, NSW 2087 (AU). (74) Agents: CORBETT, Terence, Guy et al.; Davies & Collison, 1 Little Collins Street, Melbourne, VIC 3000 (AU). (81) Designated States: AT, AU, BB, BG, BJ (OAPI patent), BR, CF (OAPI patent), CG (OAPI patent), CH, CM (OAPI pa-tent), DE, DK, FI, GA (OAPI patent), GB, HU, JP, KP, KR, LK, LU, MC, MG, ML (OAPI patent), MR (OAPI pa-tent), MW, NL, NO, RO, SD, SE, SN (OAPI patent), SU, TD (OAPI patent), TG (OAPI patent), US. Published With international search report.</p>
<p>(54) Title: RIBOZYMES</p> <p style="text-align: center;">AA G C*G AGXAGUC N₁*N₂ (I)</p>		
<p>(57) Abstract</p> <p>Synthetic single stranded RNA molecule(s) and derivatives thereof possessing highly specific endoribonuclease ac-tivity, hereinafter defined as ribozyme. The ribozyme cleaves target RNA sequence by hybridizing partially to the target RNA. It comprises of a hybridizing region and a catalytic region. The hybridizing region contains one or more arms of single stranded RNA. The preferred ribozyme has a catalytic region consisting of the sequence (I), in a loop structure where * represents base pairing between complementary nucleotides, X is any ribonucleotide and N₁ and N₂ represent ribonucleo-tides of complementary sequences along at least part of the length. The ribozyme cleaves target RNA after the triplet XUY, where X is any ribonucleotide, U is uracil and Y is adenine, cytosine or uracil. Methods for inactivating RNA (in vitro or in vivo), methods of preparing the ribozymes, and compositions containing the ribozymes are also described.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	ML	Mali
AU	Australia	GA	Gabon	MR	Mauritania
BB	Barbados	GB	United Kingdom	MW	Malawi
BE	Belgium	HU	Hungary	NL	Netherlands
BG	Bulgaria	IT	Italy	NO	Norway
BJ	Benin	JP	Japan	RO	Romania
BR	Brazil	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	LI	Liechtenstein	SN	Senegal
CH	Switzerland	LK	Sri Lanka	SU	Soviet Union
CM	Cameroon	LU	Luxembourg	TD	Chad
DE	Germany, Federal Republic of	MC	Monaco	TG	Togo
DK	Denmark	MG	Madagascar	US	United States of America
FI	Finland				

RIBOZYMES

The present invention relates to a class of synthetic RNA molecules and derivatives thereof, hereinafter referred to as ribozymes, which possess highly specific endoribonuclease activity.

5 A number of naturally occurring RNA molecules such as avocardo sunblotch viroid (ASBV), the satellite RNAs of tobacco ringspot virus (sTobRV) and lucerne transient streak virus (sLTSV), undergo self-catalysed cleavage. Such cleavage appears to be
10 an essential and unique part of the life cycle of these and other RNAs.

Self-catalysed RNA cleavage reactions share a common requirement for divalent metal ions and neutral or higher pH, and result in the production of
15 RNA with termini possessing 5' hydroxyl and 2',3' cyclic phosphate groups (Prody et al., Science 231: 1577-1580 (1986) and Buzayan et al., Virology 151: 186-199 (1986)). The cleavage reactions are
• catalysed by the RNAs themselves, presumably as a

result of conformation bringing reactive groups into close proximity. The sites of self-catalysed cleavage in naturally occurring RNAs are located within highly conserved regions of RNA secondary structure (Buzayan et al., Proc. Natl. Acad. Sci. USA 83: 8859-8862 (1986) and Forster, A.C. and Symons, R.H. Cell 50: 9-16 (1987)).

Experiments we have carried out on the satellite RNAs of tobacco ringspot virus (sTobRV) have led to the design of novel endoribonucleases (hereinafter referred to as "ribozymes"), that is, enzymes comprised of RNA which catalyse specific cleavage of RNA target molecules.

The term ribozyme as used in the specification refers to molecules comprised wholly of RNA or derivatives thereof.

The ribozymes of the present invention are distinct from the RNA endoribonuclease which occurs naturally in Tetrahymena Thermophila (known as the IVS, or L-19 IVS RNA) and which has been extensively described by Thomas Cech and collaborators (Zaug, A.J. et al, Science (1984) 224: 574-578; Zaug, A.J. and Cech, T.R., Science (1986) 231: 470-475; Zaug, A.J., et al, Nature (1986) 324: 429-433; published International patent application No. WO 88/04300 by University Patents Inc.). The Cech endoribonuclease has an eight base pair active site which hybridizes to a target RNA sequence whereafter cleavage of the target RNA takes place, with a requirement for free guanosine or guanosine derivatives. The fragments which arise from cleavage contain terminal 5' phosphate and 3' hydroxyl groups. The limited number of nucleotides available for hybridization to an RNA substrate limits the effectiveness or efficiency of

the Cech endoribonuclease as in general, oligonucleotides comprising less than twelve nucleotides hybridize poorly to target sequences. It also appears that a number of nucleotides in the active site of the Cech endoribonuclease may need to be conserved for efficient endoribonuclease activity. This restricts the number of permutations of active site sequences which can be engineered to effect hybridization to target sequences, thereby restricting the range of RNA target sequences cleavable by the Cech endoribonuclease. The Cech endoribonuclease also modifies RNA by adding a free guanosine nucleotide to the 5' end of cleaved RNA.


In contrast, the ribozymes of the present invention hybridize efficiently to a wide range of target RNA sequences, and do not modify the cleaved target RNA.

The ribozymes of the present invention comprise a hybridizing region which is complementary in nucleotide sequence to at least part of a target RNA, and a catalytic region which is adapted to cleave the target RNA. The hybridizing region contains 9 or more nucleotides.

In a preferred aspect the ribozymes of the present invention have a hybridizing region comprising one or more arms formed of single stranded RNA and having a sequence complementary to at least part of a target RNA, said one or more arms being associated with a catalytic region capable of cleaving said target RNA; and where the hybridizing region comprises a single arm of RNA said arm contains at least 9 nucleotides, and where the hybridizing region comprises 2 or more arms of RNA, the sum of nucleotides in said arms is greater than 9

nucleotides.

In one embodiment of the invention, there is provided a ribozyme of the formula 1:



- wherein; X represents any ribonucleotide and each X
 15 residue may be the same or different;
 the sum of n and n' is greater than 6 and n
 and n' may be the same or different;
 an (*) represents a base pair between
 complementary ribonucleotides;
 20 X' and X'' represent oligoribonucleotides of
 complementary sequence along at least part
 of their length to allow base pairing
 between the oligoribonucleotides, or X' and
 X'' together form a single RNA sequence
 25 wherein at least part of said sequence
 comprises a stem formed by base pairing
 between complementary nucleotides; and
 optionally, an additional nucleotide
 selected from one of A, G, C or U may be
 30 inserted after ¹A in formula (1).


Region (I) of formula (1) represents the arms or flanking sequences of a ribozyme which hybridize to respective portions of a target RNA sequence. The arms may hybridize along the full length of the

target RNA or part thereof. An RNA catalytic region is depicted at region (II) of formula 1. The catalytic region may contain one or more additional nucleotides which do not adversely effect catalytic activity. Such additions could be readily tested for ribozyme activity without undue experimentation following the teachings of the specification. The catalytic region may also form part of the hybridizing region.

10 The oligoribonucleotides X' and X" may comprise up to 5,000 or more nucleotides.

According to a further specific embodiment of the present invention there is provided a ribozyme of the formula 2:

15



wherein; X represents any ribonucleotide and each X residue may be the same or different;

30 an (*) represents a base pair between complementary ribonucleotides;

n and n' are as previously defined;

m and m' are 1 or more and may be the same or different;

B represents a bond, a base pair, a ribonucleotide, or an oligoribonucleotide containing at least 2 ribonucleotides;

and optionally, an additional nucleotide
5 selected from any one of A, G, C or U may be inserted after ¹A in formula (2).

The ribozymes of the present invention may be prepared by methods known per se in the art for the synthesis of RNA molecules. (For example, according
10 to recommended protocols of Promega, Madison, WI, USA). In particular, the ribozymes of the invention may be prepared from a corresponding DNA sequence (DNA which on transcription yields a ribozyme, and which may be synthesized according to methods known
15 per se in the art for the synthesis of DNA) operably linked to an RNA polymerase promoter such as a promoter for T7 RNA polymerase or SP6 RNA polymerase. A DNA sequence corresponding to a ribozyme of the present invention may be ligated into
20 a DNA transfer vector, such as plasmid or bacteriophage DNA. Where the transfer vector contains an RNA polymerase promoter operably linked to DNA corresponding to a ribozyme, the ribozyme may be conveniently produced upon incubation with an RNA
25 polymerase. Ribozymes may, therefore, be produced in vitro by incubation of RNA polymerase with an RNA polymerase promoter operably linked to DNA corresponding to a ribozyme, in the presence of ribonucleotides. In vivo, prokaryotic or eukaryotic
30 cells (including mammalian and plant cells) may be transfected with an appropriate transfer vector containing genetic material corresponding to a ribozyme in accordance with the present invention, operably linked to an RNA polymerase promoter such

that the ribozyme is transcribed in the host cell. Transfer vectors may be bacterial plasmids or viral RNA or DNA. Nucleotide sequences corresponding to ribozymes are generally placed under the control of strong promoters such as, for example, the lac, SV40 late, SV40 early, metallothionin, or λ promoters. Ribozymes may be directly transcribed in-vivo from a transfer vector, or alternatively, may be transcribed as part of a larger RNA molecule. For example, DNA corresponding to ribozyme sequences may be ligated into the 3' end of a carrier gene, for example, after a translation stop signal. Larger RNA molecules may help to stabilize the ribozyme molecules against nuclease digestion within cells. On translation the carrier gene may give rise to a protein, whose presence can be directly assayed, for example, by enzymatic reaction. The carrier gene may, for example, encode an enzyme.

In a further aspect of the invention, there is provided a DNA transfer vector which contains a DNA sequence corresponding to a ribozyme operably linked to a promoter to provide transcription of the ribozyme.

In one preferred method of producing a ribozyme, two synthetic oligonucleotides of complementary sequence are prepared by standard procedures (for example, using an Applied Biosystems Model 380A DNA Synthesizer (Applied Biosystems Inc., Foster City, California 94404)), and hybridized together. One of the oligonucleotides encodes a desired ribozyme. The respective ends of the hybridized oligonucleotides correspond to different restriction enzyme sites, say EcoRI at one end and PstI at the other end. After cleavage with appropriate restriction enzymes (EcoRI

and PstI in the above example), the double stranded DNA fragment may be cloned into a transfer vector. Where the plasmid vector contains an RNA polymerase promoter upstream from the DNA sequence corresponding
5 to a ribozyme of the present invention, RNA transcripts corresponding to the ribozyme can be conveniently prepared either in-vitro or in-vivo. Where the ribozyme is comprised of two halves held together by base-pairing between complementary
10 nucleotides, each half of the ribozyme may be produced according to the above methods, and the halves incubated together to form the ribozyme.

The preferred ribozymes of the present invention cleave target RNA which contains the sequence X^OUY ,
15 where X^O is any ribonucleotide, U is uracil and Y is adenine, cytosine or uracil. X^OU forms part of a base pair flanking region and Y is not base paired. Preferably, but by no means exclusively, X^O is guanidine, and X^OUY is GUC or GUA. Any RNA
20 molecule containing these sequences can be cleaved with the ribozymes of the present invention. Once the sequence of an RNA transcript containing the sequence X^OUY has been determined, the arms of the ribozyme sequence can be synthesised to be
25 complementary to, and thus hybridizable to, the RNA on the target sequence flanking the X^OUY sequence. On hybridization of the arms of the ribozyme to the target RNA sequence flanking the X^OUY sequence, the catalytic region of the ribozyme cleaves the target
30 RNA within the X^OUY sequence. RNA cleavage is facilitated in the presence of magnesium or other divalent cation at a pH of approximately 8.0.

Accordingly, the preferred ribozymes of the present invention can be engineered to cleave any RNA

whose sequence is known. The high frequency of the residues cleaved by the ribozymes in RNA (1:64 for GUC in an RNA with random and equal frequency of base distribution) means that a number of potential sites
5 for ribozyme cleavage can be confidently predicted in any given target RNA.

According to another aspect of the present invention there is provided a method for the inactivation of a target RNA sequence, which
10 comprises reacting said target RNA sequence with a ribozyme of the present invention.

In-vivo, that is, within the cell or cells of an organism, a transfer vector such as a bacterial plasmid or viral RNA or DNA, encoding one or more
15 ribozymes, may be transfected into cells e.g. (Llewellyn et al., J. Mol. Biol. (1987) 195: 115-123; Hanahan et al. J. Mol. Biol (1983) 166). Once inside the cell, the transfer vector may replicate, and be transcribed by cellular polymerases
20 to produce ribozyme RNAs which then inactivate a desired target RNA. Alternatively, a transfer vector containing one or more ribozyme sequences may be transfected into cells or introduced into cells by way of micromanipulation techniques such as
25 microinjection, such that the transfer vector or a part thereof becomes integrated into the genome of the host cell. Transcription of the integrated genetic material gives rise to ribozymes, which act to inactivate a desired target RNA.

30 The ribozymes of the present invention have extensive therapeutic and biological applications. For example, disease causing viruses in man and animals may be inactivated by administering to a subject infected with a virus, a ribozyme in

accordance with the present invention adapted to hybridize to and cleave RNA transcripts of the virus. Such ribozymes may be delivered by parenteral or other means of administration. Alternatively, a
5 subject infected with a disease causing virus may be administered a non-virulent virus such as vaccinia or adenovirus which has been genetically engineered to contain DNA corresponding to a ribozyme operably linked to an RNA promoter, such that the ribozyme is
10 transcribed in the cells of the host animal, transfected with the engineered virus, to effect cleavage and/or inactivation of the target RNA transcript of the disease causing virus. The ribozymes of the present invention have particular
15 application to viral diseases caused for example, by the herpes simplex virus (HSV) or the AIDS virus (HIV).

The ribozymes of the present invention also have particular application to the inactivation of RNA
20 transcripts in bacteria and other prokaryotic cells, plants and animals. In bacteria, RNA transcripts of, for example, bacteriophage, which cause bacterial cell death, may be inactivated by transfecting a cell with a DNA transfer vector which is capable of producing a
25 ribozyme in accordance with the present invention which inactivates the phage DNA. Alternatively, the ribozyme itself may be added to and taken up by the bacterial cell to effect cleavage of the phage RNA.

RNA transcripts in plants may be inactivated
30 using ribozymes encoded by a transfer vector such as the Ti plasmid of Agrobacterium tumefaciens. When such vectors are transfected into a plant cell, the ribozymes are produced under the action of RNA polymerase and may effect cleavage of a specific

target RNA sequence. Accordingly, plant viruses whose RNA sequence are known, or the RNA transcripts of plant genes, may be inactivated using ribozymes.

Endogenous gene transcripts in plants, animals or
5 other cell types may be inactivated using the ribozymes of the present invention. Accordingly, undesirable phenotypes or characteristics may be modulated. It may, for example, be possible using the ribozymes of the present invention to remove
10 stones from fruit or treat hereditary diseases in humans which are caused by the production of a deleterious protein, or over production of a particular protein.

The present invention will now be illustrated by
15 way of non-limiting example only, with reference to the following non-limiting Examples, and Figures.

In the Figures:

FIGURE 1 shows RNA self cleavage sites of wild type and mutated RNAs; and an electrophoretic profile
20 showing self-catalysed RNA cleavage products.

(a) Summarises the conserved structures associated with naturally-occurring RNA cleavage sites in ASBV, newt satellite DNA transcripts and the satellite RNAs of sTobRV, LTSV, velvet tobacco mottle
25 virus, solanum nodiflorum mottle virus and subterranean clover mottle virus. Nucleotide sequences which are conserved between these structures are shown, while others are represented as X. Base-pairing is represented by "*" and the site
30 for RNA cleavage is arrowed.

(b) Shows the conserved nucleotide sequence associated with the cleavage of the (+) strand of sTobRV RNA. The cleavage site is arrowed.

(c) An in vitro mutant of sTobRV containing an

insertion of eight nucleotides (shown boxed) together with a flanking duplication of three nucleotides (UGU residues 7 to 9) is shown.

(d) Sub-cloned HaeIII fragments of wild type sTobRV and the D-51 in vitro mutant were each transcribed in both the (+) and (-) orientations and radiolabelled transcripts fractionated by polyacrylamide gel electrophoresis. The positions of uncleaved 159 and 170 base transcripts from the wild type (WT) and mutant (D-51) sequences are arrowed; sizes of cleavage products are shown.

FIGURE 2 shows the nucleotide sequence of a ribozyme and the products of ribozyme cleavage separated by gel electrophoresis.

(a) The inserted nucleotides in the D-51 mutant (FIG. 1c) contains a BamHI restriction endonuclease site. BamHI was used to split the mutant DNA, and the two sequences were sub-cloned and transcribed separately in vitro. The RNA transcripts are shown schematically, with potential base-pairings between the RNAs indicated "***". The fragment containing the arrowed site for cleavage is referred to as S-RNA, the fragment containing the ribozyme is designated Rz-RNA.

(b) [^{32}P]-Rz-RNA (101 bases) was incubated alone (lane 1), and with unlabelled S-RNA (lane 2). [^{32}P]-S-RNA was incubated alone (lane 3), and with unlabelled and ^{32}P labelled Rz-RNAs (lanes 4 & 5 respectively).

FIGURE 3 shows a schematic model of a ribozyme according to one embodiment of the present invention. Region A represents the cleavage sequence within the target RNA. Region B represents a catalytic region, and regions C represent the arms of

the ribozyme.

FIGURE 4 shows the design of ribozymes targeted against the CAT (chloramphenicol acetyl transferase) gene transcript. Ribozymes, termed RzCAT-1, 2 and 3, were targeted against three sites within an 835 base in vitro transcript of the CAT gene. The relative locations of the cleavage sites on the transcript are shown schematically with the flanking bases numbered (a). The three ribozyme sequences are shown ((b) to (d)) with their target sequences. Amino-acid sequences of the CAT gene are numbered and the predicted sites for RNA cleavage arrowed. RzCAT-1 and 3 contain 24 base sequences derived from (+) strand sTobRV (region B, FIG. 3), while RzCAT-2 contains a single U-A change in this region.

FIGURE 5 shows the results of CAT RNA cleavage with ribozymes RzCAT-1 to 3.

(a) The [32 P]-CAT RNAs were gel fractionated after incubation alone (-) or with with one of the three ribozymes, RzCAT-1 to 3 (lanes 1, 2 & 3 respectively). The location of full-length transcript is shown arrowed.

(b) 5' Terminal base analysis. The 3' fragments produced by ribozyme cleavage of CAT mRNA were [5'- 32 P]-kinased, gel purified, subjected to complete nuclease digestion and the released terminal residues fractionated by pH 3.5 polyacrylamide gel electrophoresis. The 5' terminal nucleotides, determined by reference to markers (lane M), were A, U and G for the fragments produced by RzCAT-1 to 3 (lanes 1, 2 and 3 respectively).

FIGURE 6 shows a time course of ribozyme (RzCAT-1) catalytic activity against CAT RNA. The amounts of the 139 nucleotide cleavage product were

quantified and plotted. The inset shows the accumulation of the 139 base fragment with time, after polyacrylamide gel electrophoresis.

FIGURE 7 shows the relative rates of cleavage of CAT RNA under different temperature conditions. Substrate RNA is represented by a solid line. In each case cleavage product is represented by a broken line.

FIGURE 8 depicts three ribozymes (corresponding to RzCAT-2) having arms or flanking sequence of varying length.

FIGURE 9 shows the scheme for production of a ribozyme comprising catalytic anti-sense RNA containing each of the cleavage domains RzCAT-1 to 3.

FIGURE 10 shows ribozymes hybridizing to target sequences containing the GUA (10a) and GUU (10b) motifs in CAT mRNA.

FIGURE 11 shows sites for self catalysed RNA cleavage in citrus exocortis viroid (CEV) RNA and its complement.

FIGURE 12 shows the ribozyme RzCEV25x(+) hybridized to target CEV RNA(a), and a gel electrophoretic profile of (+)CEV RNA and complementary (-) CEV RNA incubated with RzCEV25x(+) (b, lanes 1 and 2 respectively. Cleavage product is arrowed).

FIGURE 13 shows the ribozyme RzCAT-2 hybridizing to its target sequence (a) and ribozyme RzSCMoV(b). The catalytic domain in each ribozyme is boxed. Differences in the catalytic region of RzSCMoV, when compared with RzCAT-2 are marked.

FIGURE 14 shows the ribozyme RzCEV-2 hybridizing to a target sequence in citrus exocortis viroid (CEV) RNA. The cleavage site corresponds to nucleotide-336

in the CEV RNA sequence. The alteration in nucleotide sequence in the catalytic domain, when compared with the catalytic domain of sTobRV, is circled (a). Fig 14(b) shows an electrophoretic
5 profile of a control [(-) strand of CEV] RNA, lane 7 and the (+) strand of CEV RNA lane 8, after incubation with RzCEV2.

FIGURE 15 shows the ribozyme RzCAT-2 (a) compared with ribozyme RzCAT-2B (b). Catalytic domains are
10 boxed. Changes in the catalytic domain of RzCAT-2B compared with RzCAT-2 are also boxed;

FIGURE 16 shows a map of plasmid pJ35SN; and

FIGURE 17 is a graphic presentation of the average of four experiments on the inhibition of CAT
15 expression in plants (tobacco protoplasts).

The following Examples are given to illustrate the present invention and are not to be construed as limiting the present invention.

Reactions and manipulations involving DNA, such
20 as ligations, restriction enzyme digestions, bacterial transformation, DNA sequencing etc. were carried out according to standard techniques, such as those described by Maniatis et al (Molecular Cloning, Cold Spring Harbour, 1982). Manipulations involving
25 RNA were also carried out according to standard techniques, such as those described by Uhlenbeck (Nature 328, 596-600 (1987)) and Haseloff and Gerlach (Nature 334, 585-591 (1988)).

30 **EXAMPLE 1**

Self-Catalyzed Cleavage of Mutated sTobRV RNA:

A consensus of the domains associated with naturally-occurring RNA cleavage sites in ASBV, newt satellite DNA transcripts and the satellite RNAs of

sTobRV, LTSV, velvet tobacco mottle virus (VMoV), solanum nodiflorum mottle virus (SNMV) and subterranean clover mottle virus (SCMoV) as shown in Figure 1a. Nucleotide sequences which are conserved
5 between these structures are shown, while non-conserved sequences are represented as X. An extra U is positioned after residue ¹A in LTSV(+) strand.

The domain associated with the self-catalysed
10 cleavage of the (+) strand of sTobRV was studied to ascertain the enzymic substrate activity within this domain. First, cloned sTobRV cDNAs were mutagenised using an oligonucleotide linker (BamH1) insertion protocol.

15 Construction of a Vector for in vitro Expression of sTobRV:

A 160 bp Taq 1 - Spe 1 fragment of sTobRV cDNA was isolated from pSP653 (Gerlach et al. 1985, Virology 151: 172-185) and ligated to Acc 1 - Spe 1
20 cut, phosphatase-treated pGEM 4 to reform the Acc 1 site. A resulting clone was linearized with Acc 1, phosphatase-treated and a 359 bp Taq 1 fragment of the sTobRV cDNA was inserted. The resulting clones were screened for the presence of a circularly
25 permuted 520 bp sTobRV cDNA sequence containing the terminally redundant residues 277 to 81 (pTTS). The sTobRV sequence is flanked by promoters for T7 and SP6 RNA polymerases, and in vitro transcription gave rise to RNAs of (+) or (-) orientation which
30 contained two sites for self-cleavage.

In vitro Mutagenesis:

The plasmid pTTS (50 ug) was linearized with BamH 1, treated with S1 nuclease and religated, to remove a unique BamH 1 site. The resulting construction,

pTTS-B, was treated with 2×10^{-4} units DNase I in 20mM Tris-HCl pH 7.0, 15 mM $MnCl_2$ for 10 mins. at 37°C. The resulting linear DNAs were trimmed and/or end-filled using T4 DNA Polymerase, and purified by 5 0.7% LGT agarose gel electrophoresis and extraction. Kinased BamH I linker sequences (CGGATCCG) were ligated to the linearized plasmid overnight at room temperature in the presence of 5% polyethylene glycol. Subsequently, the reactions were BamH I 10 digested, and the linear plasmid DNAs repurified by 0.7% LGT agarose gel electrophoresis (this was found necessary to remove last traces of circular plasmid, together with unligated linkers). Plasmids were recircularized using T4 DNA ligase and transformed 15 into E. coli DH-1. Colonies (greater than 1000) were scraped from agar plates, grown in liquid culture to saturation and a mixed population of plasmid DNAs prepared. The mixed sTobRV cDNA inserts were excised by restriction enzyme digestion at flanking EcoRI and 20 PstI sites, purified by 1% LGT agarose gel electrophoresis, and sub-cloned into EcoRI - PstI cut, phosphates-treated pGEM 4. The resulting transformants were again pooled, grown in liquid culture and plasmid DNA prepared. The plasmid DNAs 25 were treated with BamHI, to cleave only those plasmids containing a BamHI linker sequence, and the linear forms were again purified by two rounds of 0.7% LGT agarose gel electrophoresis, recircularized with T4 DNA ligase, and transformed into E. coli 30 DH-1. Individual transformants were screened for the approximate position of the inserted BamHI linker within the sTobRV sequence by restriction enzyme digestion, sub-cloned into M13 mp19 and sequenced via the dideoxynucleotide chain termination technique.

A library of sTobRV mutants resulted, and nucleotide sequence analysis showed that each mutant contained an inserted BamHI linker sequence (CGGATCCG) together with flanking duplicated or
5 deleted sTobRV sequences. The mutants were transcribed in vitro and the RNAs assayed for their ability to undergo cleavage. From these experiments, a 52-nucleotide sequence was identified as containing both the substrate and cleavage portions of sTobRV
10 RNA. This 52-nucleotide sequence, depicted in Figure 1b, contained the domain of conserved sequence required for self-cleavage of other RNAs (Figure 1a). One mutant, designated D-51, contained an eight nucleotide BamHI linker sequence inserted between
15 three duplicated sTobRV nucleotides numbered 7 to 9. This mutant underwent self-catalysed RNA cleavage.

97 and 108 base-pair HaeIII fragments containing the 52-nucleotide cleavage sequence of the wild type and D-51 RNAs (as shown in Figures 1b and 1c) were
20 excised from sequenced plasmid clones. The fragments were ligated into the SmaI site of pGEM4 and screened to obtain both orientations of the insert. The plasmids were linearised using EcoRI and (+) and (-) strand RNAs of lengths 159 and 170 bases were
25 transcribed using 200 units/ml T7 RNA polymerase in 50 mM Tris-HCl, pH 7.5, 10 mM NaCl, 6 mM MgCl₂, 2 mM spermidine, 1000 units/ml RNasin, 500 µM ATP, CTP and GTP with 200 µM [α ³²P] UTP. RNAs were fractionated by electrophoresis on a 10%
30 polyacrylamide, 7 molar urea, 25% formamide gel, and autoradiographed.

As shown in Figure 1d, no cleavage of the (-) strand RNA transcripts was observed. This was as expected, as the (-) strand did not contain a

self-catalysed cleavage site. With the (+) strands of both the wild type and D-51 sequences, cleavage took place, with cleavage of the D-51 RNA being somewhat less efficient than that of the wild type (Figure 1d). This experiment indicates that the single stranded loop region at the right-hand side of the 52-nucleotide sequence involved in the self-catalysed cleavage of RNA is non essential.

Separation of Enzymic and Substrate Activities:

- 10 Using the BamHI restriction endonuclease site inserted into D-51, the flanking HaeIII-BamHI and BamHI-HaeIII fragments were obtained and each was sub-cloned into an E. coli plasmid suitable for in-vitro transcription. This led to the elimination
- 15 of the mutated single-stranded loop from the self-cleavage domain, splitting the region into two RNA segments (Figure 2a). The smaller HaeIII-BamHI fragment contained nucleotides 321 to 9, including the actual site of cleavage and was termed the S
- 20 fragment. The BamHI-HaeIII fragment containing sTobRV nucleotides 7 to 48 was termed the ribozyme or Rz fragment. The E. Coli plasmids used for in-vitro transcription were pGEM3 and pGEM4 (Promega, Madison, WI, U.S.A.). These expression plasmids contain:
- 25 (a) an origin of replication;
(b) selectable drug resistance (Amp^r) gene;
(c) a multiple cloning site flanked by RNA polymerase promoters which can be used for in vitro production of transcripts.
- 30

T7 DNA polymerase treated, KpnI digested Rz-pGEM3 and XbaI digested S-pGEM4 were transcribed using SP6 RNA polymerase under the same conditions as set out above.

As shown in Figure 2, both the S and Rz-RNAs showed no significant degradation when incubated alone (Figures 2b, lanes 1 and 3) under conditions suitable for highly efficient self-cleavage (50°C, 20mM MgCl₂, pH 8.0). The labelled Rz-RNA also appeared unaltered after incubation with the S-RNA (Figure 2b, lanes 2 and 5). However, when the S-RNA was mixed with the Rz-RNA, efficient cleavage of the S-RNA occurred (Figure 2b, lanes 4 and 5) producing two fragments. The product sizes were consistent with cleavage of the S-RNA (84 bases) at the normal site between nucleotides #359 and #1, to give 5' and 3' proximal fragments of 67 and 17 nucleotides, respectively. This shows that the S-RNA acted as a substrate for ribonucleolytic cleavage by the Rz-RNA, which acted in a catalytic fashion.

A model of a ribozyme based on the catalytic region of sTobRV RNA is shown in Figure 3. The ribozyme has two arms or flanking sequences of single stranded RNA shown at C, hybridizing to complementary sequences on a substrate RNA, i.e. RNA to be cleaved. Each flanking sequence shown at C, contains 8 ribonucleotides. The number of nucleotides contained in region C is not critical. Sufficient nucleotides must, however, be present to allow the ribozyme to hybridize to a target RNA. Four nucleotides in each region C appears to be the minimum number for hybridization.

The catalytic region B contains sequences which are highly conserved in naturally-occurring cleavage domains (see Figure 1a). From a comparison with cleavage domains of the known sequences, the length of the base pair stem II is unimportant, as is the presence of an associated loop at one end thereof.

The cleavage site within the target RNA is depicted at A (in Figure 3) as GUC. On the basis of our experiments (not shown), and others by Koizumi (FEBS LETT 288; 228-230 (1988); and FEBS LETT 239; 285-288 (1988)) on the cleavage sites in naturally occurring RNAs, the sequences GUA, GUC, CUC, AUC and UUC also act as cleavage sites within RNA.

EXAMPLE 2

10 Demonstration of the Design, Synthesis and Activity of Ribozymes with New and Highly Specific Endoribonuclease Activity:

As an illustration of this invention, three ribozymes have been designed, which are targeted against the transcript of a commonly used indicator gene derived from bacteria, Tn9 Chloramphenicol Acetyl Transferase (CAT), which can provide antibiotic resistance in bacteria, plants and animals and can be easily assayed. These ribozymes, designated RzCAT-1 to 3 correspond to potential GUC cleavage sites in CAT RNA at positions 139-140, 494-495 and 662-663 respectively. The sequences of these ribozymes are depicted in Figure 4. In each case, the flanking sequences of the ribozyme which hybridize to the target CAT RNA, were 8 nucleotides in length. The catalytic region was chosen to correspond to that of sTobV RNA, shown in Figure 3.

The CAT gene was obtained from pCM4 and sub-cloned as a BamHI fragment into pGEM-32 (from Promega, Madison, WI, U.S.A.). This plasmid was linearised with HindIII and CAT gene transcripts were obtained using T7 RNA polymerase with 220uM [α -³²P]UTP. Ribozym sequences were synthesised

as oligodeoxynucleotides, Rz CAT-1, 2 and 3, respectively. They were kinased, ligated with phosphatased treated, EcoRI-PstI cut pGEM4 and incubated with the Klenow fragment of DNA polymerase 5 1 before bacterial transformation. EcoRI linearised plasmids were transcribed with T7 RNA polymerase to produce ribozyme RNAs. Ribozymes were incubated with CAT transcript in 50 mM Tris-HCl pH 8.0, 20 mM $MgCl_2$ at 50°C for 60 min, and the products 10 fractionated by 5% polyacrylamide 7M urea, 25% formamide gel electrophoresis prior to autoradiography.

When the 840-nucleotide CAT transcript was incubated with any one of the three ribozymes, 15 efficient and highly sequence-specific cleavage occurred (Figure 5) producing 2 RNA fragments in each reaction. The fragment sizes were consistent with the predicted sites for cleavage (i.e. 139 and 696, 494 and 341, 662 and 173 base fragments were the 5' 20 and 3' products from RzCAT-1 to 3 catalysed cleavage respectively). The conditions required for these ribozyme-catalysed cleavages were similar to those observed for naturally occurring cleavage reactions (Foster, A.C. and Symons, R.H., Cell 49: 211-220 25 (1987) and Foster, A.C. and Symons, R.H. Cell 50: 9-16 (1987)), with more efficient cleavage occurring at elevated pH, temperature and divalent cation concentrations (data not shown). When present in molar excess, the three ribozymes catalysed almost 30 complete cleavage of the CAT RNA substrate after 60 min. in 50 mM Tris HCl, pH 8.0, 20 mM $MgCl_2$ at 50°C. Under similar conditions with 0.1 μM substrate and 3 μM ribozymes, the $T_{1/2}$ of CAT mRNA substrat was 3.5, 3.5 and 2.5 min. in the

presence of RzCAT-1 to 3 respectively. The ribozyme sequences were inactive against the complement of the substrate RNA (i.e. the (+) strand), and in the form of oligodeoxyribonucleotides (data not shown). The

5 3' terminal cleavage fragments from each ribozyme catalysed reaction were isolated and 5' ^{32}P -kinased (50 mM TrisHCl pH 9, 10 mM MgCl_2 , 10 mM DTT with 50 uCi γ - ^{32}P ATP and 5 units T4 polynucleotide kinase for 30 min. at 37°C.).

10 Efficient kinasing of the fragments indicated that they possessed 5' terminal hydroxy groups, similar to those produced in naturally occurring cleavage reactions.

The terminal nucleotide of the fragments produced

15 by cleavage of the CAT sequences by RzCAT-1 to 3 were determined. Briefly, radiolabelled fragments were purified on a 5% polyacrylamide gel and digested with an equal volume of 500 units/ml RNase T1, 25 units/ml RNase T2 and 0.125 mg/ml RNaseA in 50 mM ammonium

20 acetate pH 4.5 for 120 min. at 37°C. The products were fractionated on a 20% polyacrylamide gel containing 25 mM sodium citrate, pH 3.5 and 7 molar urea. Figure 5b shows that the cleavage of the CAT sequences by RzCAT-1 to 3 occurs precisely before

25 nucleotides A, U and G respectively.

The terminal sequence of the CAT gene fragments were determined directly using the partial enzymatic digestion technique (Donis-Keller et al., Nucleic Acids Res. 4: 2527-2538 (1980)), using base-specific

30 partial ribonucleolytic cleavage. The sequence of the fragments confirmed that cleavage occurred at the expected locations within CAT RNA (not shown).

Enzymatic Catalysis:

To demonstrate that ribozymes cause cleavage of

the CAT mRNA substrate in a catalytic manner, each was incubated with a molar excess of substrate, under conditions which should favour both efficient cleavage and product dissociation. Figure 6 shows the results of an experiment where after 75 min. at 50°C, pH 8.0 in 20 mM MgCl₂, 10 pmoles of RzCAT-1 had catalysed specific cleavage of 163 pmoles of a truncated CAT mRNA (173 bases) substrate to give 5' and 3' fragments of 139 and 34 bases, respectively. On average, each ribozyme had participated in greater than ten cleavage events. After 75 minutes at 50°C some non-specific cleavage of RNA was noticed due to the extreme conditions, but 70% of the remaining intact RNAs (163 pmoles) had accumulated as the 139 base fragment. Similar results were obtained for RzCAT-2 and 3 (data not shown), and thus each acts as an RNA enzyme.

EXAMPLE 3

20 The Effect of Temperature on Ribozyme Activity:

The effect of reaction temperature on the in-vitro rate of ribozyme activity was examined.

A time course of reactions for ribozymes RzCAT-1 to 3 on CAT RNA substrate at 37°C and 50°C was 25 carried out.

In this experiment, reactions for each ribozyme were set up in duplicate, using reaction conditions for ribozyme cleavage set out in Example 2. One reaction was incubated at 37°C, the other at 50°C. 30 Samples were removed at time points to 90 minutes and the extent of reaction was analysed by denaturing polyacrylamide gel electrophoresis. Figure 7 shows the time course of reaction for each of the ribozymes RzCAT-1 to 3 at 37°C and 50°C. The reaction rate of

each ribozyme increases with increased reaction temperature.

The time taken for 50% ($t_{1/2}$) cleavage of CAT RNA is set out in Table 1.

5

TABLE 1

	RzCAT-1	RzCAT-2	RzCAT-3 $t_{1/2}$ (mins.)
50°C	3.5	3.5	2.5
10 37°C	55.0	70.0	65.0

As shown in Table 1, the rate of reaction of ribozymes at 37°C is approximately 20 times slower
15 than the rate of reaction at 50°C.

EXAMPLE 4

The Effects of Varying Arm Lengths of Ribozymes (or Flanking Sequence) on Ribozyme Catalytic Activity:

20 The arms or flanking sequences of a ribozyme (region (I) of formula 1) hybridize the ribozyme to a target RNA whereafter cleavage of the RNA takes place. In this experiment, the effect on cleavage rate of a target sequence by altering the extent of
25 complementarity and subsequent length of base pairing of the ribozyme arms to the target sequence was investigated.

Ribozymes were produced with 4, 8 and 12 base complementarity to the target sequence RzCAT-2 on
30 each arm (Figure 8a). The ribozymes were prepared according to the methods of Example 2. Ribozyme activity was determined by incubating the ribozyme RNA with CAT RNA as described previously.

The ribozyme having a 4 base complementarity on

each arm did not cleave the substrate RNA. The ribozyme with 8 base complementarity on each arm cleaved the CAT substrate as did the ribozyme having 12 base complementarity. Ribozymes with 12 base complementarity cleaved target RNA more efficiently, as judged by reaction rate in-vitro than did ribozymes having a lesser number of base complementarities. Even though it appears necessary to have more than four base complementarity, increasing the length of the hybridizing region of ribozymes increases their reaction rate.

In a second experiment, the reaction efficiency of a ribozyme having (a) complementarity to the entire length of the CAT transcript target RNA and (b) multiple catalytic domains, was investigated.

Four GUC target sites in CAT RNA sequences were chosen. Ribozyme catalytic domains against these sites were "inserted" into a complete anti-sense (-) sequence for the CAT transcript and catalytic activity tested.

The four sites chosen were the three specified by RzCAT-1 to 3 described previously, and a further site which may be represented as follows:

New CAT site

#192

	His	His	Ala	Val	Cys	Asp	Gly	
5'	CAU	CAU	GCC	GUC	UGU	GAU	GGC	3'

where "192" refers to amino acid 192 in the CAT polypeptide and an refers to the site of cleavage.

Oligodeoxyribonucleotides containing ribozyme catalytic domains and spanning each of these cleavage sites were used for M13 mutagenesis experiments to produce a sequence containing the entire complement of the CAT sequence but with the four ribozyme

catalytic domains inserted within it. M13 mutagenesis was performed by binding of oligonucleotides containing ribozyme insertions to single stranded M13 DNAs containing uracil, followed
5 by synthesis of complementary DNAs containing the insertion. The complementary DNAs were recovered following cloning in an appropriate Escherichia coli strain, (T.A. Kunkel 1985, Proc. Natl. Acad. Sci. U.S.A. 82: 488-492). The resultant double-stranded
10 cDNA was cloned in an in-vitro expression vector to produce ribozyme RNA using the T7 polymerase transcription system. Ribozyme activity was determined by incubation of ribozyme RNA with CAT transcript followed by gel electrophoresis of the
15 reaction mixture, after glyoxal treatment to denature nucleic acids.

Autolytic cleavage occurred at all the expected sites on the CAT transcript. Accordingly, the flanking sequences of arms or a ribozyme may extend
20 along the full length of the RNA transcript which is to be cleaved.

Figure 9 schematically shows the production of catalytic anti-sense RNA containing each of the four ribozymes. The catalytic anti-sense RNA contains
25 approximately 900 bases.

Under the above reaction conditions, the ribozyme and target sequences form high molecular weight complexes presumably by extensive base pairing. A strong denaturing treatment such as glyoxal treatment
30 is required to resolve reaction products during electrophoresis.

EXAMPLE 5**Target Sequences for Ribozyme Cleavage:**

The GUA motif in mRNA was tested to see whether a ribozyme would effect RNA cleavage at this sequence.

5 A specific site in CAT mRNA, including the GUA motif (Figure 10a) was chosen and an appropriate ribozyme sequence was prepared and tested for activity. The ribozyme contained arms of 8 ribonucleotides.

10 Synthetic oligonucleotides corresponding to the ribozyme of Figure 10 were prepared according to Example 2 and double-stranded cDNA cloned into an in-vitro expression vector (pGEM 4, see above) in E. coli in order to produce ribozyme RNA using the T7
15 polymerase transcription system. Ribozyme activity was determined by incubation of ribozyme RNA with CAT mRNA, followed by gel electrophoresis of the reaction mixture as previously described.

The ribozyme effected cleavage at the GUA target
20 site (not shown). Accordingly, the motif GUA in RNA is a substrate for ribozymes of the present invention. This was not completely unexpected since one naturally occurring cleavage site in the satellite RNA of lucerne transient streak virus requires
25 recognition of a GUA site.

Similarly, a GUU motif in the CAT RNA target sequence was tested with an appropriate ribozyme (See Figure 10b), and cleavage was effected.

30 EXAMPLE 6**Ribozyme Cleavage of Viral RNA:**

Viroid RNA, in the form of citrus exocortis viroid RNA, was cleaved using a ribozym of the present invention.

Two GUC target sites were chosen in citrus exocortis viroid (CEV) RNA. One site in the complementary strand sequence was also chosen. Ribozymes were prepared against all of these sites and tested for activity. The ribozymes were designated CEV9x(+), CEV9x(-) and CEV25x(+). Figure 11 shows the three cleavage sites in CEV RNA for each of these ribozymes.

Ribozymes were prepared according to previous methods. Ribozyme RzCEV25x(+) is shown in Figure 12. This ribozyme cleaves the GUC motif at nucleotide 116 of CEV RNA.

Figure 12(b) shows cleavage of CEV RNA with ribozyme RzCEV9x(+). No cleavage is observed with ribozyme RzCEV9x(-).

This experiment indicates that ribozymes are active against target RNA sequences from diverse sources. This is to be expected, as all RNAs are formed from the basic ribonucleotide building blocks of adenine, guanine, cytosine and uracil, regardless of their animal, plant or microbial origin.

EXAMPLE 7

Examples of Ribozymes Having Variable Catalytic Domains

A ribozyme targeted against a CAT-2 site was prepared using the catalytic domain sequence from the satellite RNA of subterranean clover mottle virus (SCMoV). Twelve base complementarity of ribozyme arm flanking sequence was incorporated into the design of the ribozyme RzSCMoV. The ribozymes RzCAT-2 and RzSCMoV are shown at Figures 13a and 13b respectively. The loop region of RzSCMoV contains 5 nucleotides, having the sequence AAAUC. This is to be contrasted with the loop region of RzCAT-2 which

contains 4 nucleotides having the sequence AGAG. In addition, RzSCMoV contains a C in the catalytic region, in place of U* in RzCAT-2. The different sequences in RzSCMoV when compared with RzCAT-2 are marked.

The RzSCMoV was produced according to Example 2. RzSCMoV was active, yielding two cleavage products as expected.

In another experiment, the citrus exocortis viroid (CEV) target site at nucleotide -336 in its complementary RNA was cleaved using a ribozyme (RzCEV2) having the sequence set out in Figure 14a. The loop region designated with the letter "L" in Figure 14 comprises six nucleotides having the sequence 3'-CCTATA-5'. This is distinct from the loop region of sTobRV which comprises four nucleotides having the sequence 3'-AGAG-5'. This ribozyme cleaves target CEV complementary RNA at position -336 as shown in the electrophoretic profile of Figure 14b.

This experiment indicates that the number of nucleotides, and nucleotide sequence of the loop region is unimportant in ribozyme activity. In these experiments, the ribozyme was produced according to methods previously described in the specification.

In another experiment, the effect of base pairing in the catalytic domain (stem region) on ribozyme activity was investigated.

A modified ribozyme corresponding to RzCAT-2 but containing four extra base pairs was prepared and tested. In Figure 15a, the sequence of the ribozyme Rz CAT-2 is shown hybridized to target CAT RNA. The test ribozyme is shown at 15b, with the additional base pairs boxed. The test ribozyme had comparable

activity to that of RzCAT-2. This indicates that the base paired region of the ribozyme catalytic domain may be of variant length, without effecting catalytic activity.

5 We have observed (data not shown) that the stable in vivo form of sTobRV RNA transcripts expressed in transgenic plants is primarily circular, presumably due to ligation of 5' and 3' termini. Therefore, the use of two autolytic cleavage sites flanking a
10 sequence of interest in an in vivo RNA transcript is likely to lead to a circularized product which may have greater stability than linear transcripts. This approach appears to provide a novel method for in vivo stabilization of ribozyme sequences. This is
15 termed circularization.

EXAMPLE 8

In-vivo Activity of Ribozymes:

The in-vivo activity of ribozymes in plant cells
20 is investigated in this Example.

Experimental Protocol:

Plasmids containing anti-CAT (CAT=chloramphenicol acetyl transferase) or combined anti-CAT/ribozyme gene constructions (see below) were introduced into
25 tobacco protoplasts in the same amount and proportion relative to each other, along with another plasmid which contained a functional CAT gene construction. CAT activities were measured and compared with the base level of gene activity.


30 Materials and Methods:

(a) Electroporations and CAT assays.

These were performed as described in Llewellyn et al. J. Mol. Biol. (1987) 195:115-123. Briefly, protoplasts of Nicotiana plumbaginifolia line T5 were

(b) Gene Constructions.


The active CAT gene construction was borne on the plasmid designated pCAT7+. It was derived by fusion of a CAT gene sequence (from plasmid pCM4, see T.J. Close and R. Rodriguez, 1982, Gene 20:305-316) into the plasmid pJ35SN (derived from p35SN, W.L. Gerlach et al., 1987, Nature 328:802-805) so that the active gene construction was:




35S refers to the 35S CaMV (cauliflower mosaic virus) promoter, NOS to nopaline synthetase polyadenylation signal, T/C to transcription.

Along with 0.2 ug of pCAT7+ there were added various gene constructions in excess as described below. The gene constructions were contained within plasmids with the following designations:


- 5 pJ35SN - This vector plasmid, a map of which is shown in Figure 16, contains a 35S CaMV promoter and plant nopaline synthase 3' polyadenylation signal, which may be depicted as follows:



- pCAT7- - This contains the CAT gene sequence inserted into pJ35SN such that transcription will result in the production of the antisense CAT RNA, which may be depicted as follows:



- pCAT19- - This contains the CAT gene with four catalytic ribozyme domains included within it, (see Example 4 and Figure 9), inserted into pJ35SN such that transcription will result in the production of antisense CAT RNA containing four catalytic ribozyme domains, which may be depicted as follows:



catalytic domain insertions

Results:

- 30 The following table shows the relative CAT activities in cells 20 hours after electroporation. Activity is expressed as per cent conversion of chloramphenicol substrate in a 1 hour assay.

		ug Plasmid Electroporated			
Treatment	pCAT7+	pJ35SN	pCAT7-	pCAT19-	% Conversion
1A					0
1B					0
5 2A	0.2	18			21
2B	0.2	18			46
3A	0.2	9	9		28
3B	0.2	9	9		32
4A	0.2		18		26
10 4B	0.2		18		19
5A	0.2	9		9	19
5B	0.2			9	22
6A	0.2			18	14
6B	0.2			18	16

15 (for each treatment "A" and "B" are duplicates)

The following conclusion can be drawn from these results:

- 20 (a) The introduction of the CAT gene construction results in significant CAT activity - compare 2A,B with 1A,B. There is variability between duplicates. From the trends seen in the other samples (see "b" and "c" below) it is likely that 2A shows an abnormally low activity.
- 25 (b) Concomitant introduction of an antisense gene construction results in a decrease in the level of activity - compare 3A,B and 4A,B with 2B. The extent of the decrease is related directly to the level of the antisense gene added as plasmid -
- 30 compare 3A,B and 4A,B.
- (c) Concomitant introduction of the combined antisense/ribozyme gene construction results in a decrease in gene activity - compare 5A,B and 6A,B with 2B. Furthermore, the decrease is more

marked than for the corresponding levels of antisense gene constructions - compare 5A,B with 3A,B, and 6A,B with 4A,B.

The average results for four in-vivo experiments are shown in Figure 17. In this Figure, "control" represents treatment 2. "Antisense" represents treatment 4. "Catalytic" represents treatment 6 and "Background" represents treatment 1.

The catalytic ribozyme inhibits CAT activity an average of 47%, compared to an average of 34% for an antisense ribozyme.

The introduction of ribozyme-bearing genes into plant cells inhibits the activity of genes against which they are targeted. Furthermore, the inhibition is greater than for corresponding antisense RNA molecules.

These results show that ribozymes will be active in animal, plant or microbial cells against a range of target RNA molecules.

The mechanisms of action of the ribozymes in this Example is unclear. For example, the antisense ribozyme may irreversibly hybridize to a target RNA and catalyse phosphodiester bond cleave at one or more selected target sites along the target RNA.

Alternatively, cellular enzymes may unwind the antisense RNA from its target sequence, such that the target RNA is cleaved into two or more fragments.

EXAMPLE 9**In-vivo Activity of Ribozymes in Animal Cells:**

The activity of ribozymes in inactivating a target RNA in mammalian cells is demonstrated in this
5 Example.

Materials and Methods:

The active gene constructions encoding ribozymes were transfected into the widely available monkey kidney cell line COS1 by electroporation. In this
10 method, 3×10^6 /ml COS1 cells suspended in a buffered saline with 10% FCS (foetal calf serum), were contacted with various gene constructs and an electric discharge applied to effect electroporation of DNA into the cells. The transfected cells were
15 incubated at 37°C for 48 hours in culture medium before assay for CAT and luciferase activity.

CAT gene constructs were borne on the plasmid designated pTK CAT (Miksicek et al., Cell 46: 283-290, 1986). This plasmid was derived by the
20 introduction of a CAT gene sequence into the plasmid pSV2 such that it is under the control of the thymidine kinase promoter of the herpes simplex virus.

Gene constructs encoding ribozymes were borne on the plasmid pSV232A (De Wet et al., Molecular and
25 Cellular Biology 7: 725-737, 1987) containing the luciferase gene fused to the SV40 early promoter. DNA encoding ribozymes was ligated into the XbaI site at the 3' end of the luciferase gene according to the standard methods of Maniatis et al. (Molecular
30 Cloning, A Laboratory Manual, Cold Spring Harbour, 1982).

The following constructs were prepared using standard techniques of Maniatis et al. (Supra):
pFC58 - This plasmid vector contains DNA encoding


ribozyme RzCAT-1 fused to the 3' end of the luciferase gene in a non-functional orientation.

This may be depicted as follows:



where 232A refers to pSV232A sequences, SV40 early refers to the early promoter of SV40 and small T is DNA encoding the small T intervening sequence of SV40. This construct results in production of an RNA molecule encoding luciferase and the ribozyme RzCAT-1, the latter being in an orientation such that it would not be expected to be catalytic.

- 15 pFC4 - This plasmid is the same as pFC58 except that RzCAT-1 is replaced with RzCAT-3.
- pFC1-6 - This plasmid is the same as pFC58 except that RzCAT-1 is replaced with RzCAT-3 in the sense orientation (5'-3').
- 20 pFC20 - This plasmid is the same as pFC1-6 except that RzCAT-3 is replaced with RzCAT-2 having eight nucleotide flanking sequences.
- pFC12 - This plasmid is the same as pFC20 except that the ribozyme RzCAT-2 contains twelve
- 25 nucleotide flanking sequences.
- pFC50 - This plasmid contains the CAT gene with four catalytic ribozyme domains included within it (see Example 4 and Figure 9) in the sense orientation (5'-3'), which on
- 30 transcription gives rise to an inactive ribozyme. This plasmid may be depicted as follows:



- pFC54 - This plasmid is the same as pFC50, except that the CAT gene and the ribozyme domains are in the antisense (3'-5') active orientation.
- 5 pFC64 - This plasmid shares the SV40 promoter and polyadenylation signals with pFC50 and contains the wild type CAT gene with no inserted ribozyme domains. This gene is in an antisense orientation, and thus does not
- 10 produce CAT protein.
- pFC65 - This plasmid is the same as pFC64 except that the wild type CAT gene is in the sense (5'-3') orientation and is thus productive of CAT protein.

15 Assays:

Luciferase activity was assayed according to the methods of De Wet et al. (Supra). Briefly, COS cells were lysed 48 hours after transfection, and the cell lysate incubated with luciferin, the substrate of

20 luciferase, and luminescence detected using a scintillation counter.

CAT activity was also measured using COS cell lysates (cell lysates were divided into two, and each portion assayed either for luciferase or CAT

25 activity), according to the method of Sleight, M.J., Anal. Biochem. 156: 251-256, (1986).

In the in-vivo assays, pFC58 and pFC4 did not effect CAT activity in transfected cells. This activity was designated 100% CAT activity and 0% CAT

30 suppression. CAT activity in cells transfected with other plasmids was measured relative to pFC58.

The % of CAT suppression was measured as

39

$$100 - \left(\frac{\text{CAT}_{\text{test}}}{\text{CAT}_{\text{control}}} \right) \times 100$$

normalised to luciferase production. CAT_{test} = CAT assay result for test constructs. $\text{CAT}_{\text{control}}$ = CAT assay for control constructs (pFC4 and pFC58).

Luciferase production is an internal control for electroporation, and gives a measure of ribozyme production within each individually electroporated tissue culture plate.

Results:

Experiment (i)

μg plasmid electroporated/ 1.5×10^6 cells						
Treatment	pTKCAT	pFC58	pFC20	pFC1-6	pFC12	% CAT Suppression
15	1	5	2			56
	2	5	1	1		53
	3	5			2	40
	4	5	2			0

All treatments were carried out in duplicate and an average value given.

Experiment (ii)

20	<u>Experiment 111</u>						
	μg plasmid electroporated/ 1.5×10^6 cells						
	Treatment	PTKCAT	pFC-1-6	pFC20	pFC12	pFC4	% CAT Suppression
	5	5	2				75
	6	5		2			75
	7	5		4			62
25	8	5	1	1			70
	9	5			4		51
	10	5				2	0

Treatments 5 to 10 were carried out in duplicate and an average value given.

Experiment (iii)

<u>Experiment 11117</u>				
	μg plasmid electroporated/ 1.5×10^6 cells			
Treatment	pTKCAT	pFC1-6	pFC4	% CAT Suppression
11	5	2		66
12	5		2	0

Treatments were carried out in quintuplicate and an average value given.

Experiment (iv)

Treatment	pTKCAT	µg plasmid electrophorated				% CAT Suppression
		pFC50	pFC54	pFC64	pFC65	
13	5	2				0
14	5		2			26
15	5			2		2
16	0				2	NA

Each of treatments 13 to 16 were carried out in quadruplicate.

The sense CAT construction (treatment 16) was productive of high levels of CAT activity in its own right. Therefore % suppression is not applicable (NA).

A number of experiments were also conducted with the TK promoter of pTKCAT replaced with the human metallothionein promoter. When this CAT construct was co-transfected into COS1 cells with plasmids encoding one or more ribozymes, a marked decrease in CAT activity was observed.

The above results clearly demonstrate the in-vivo inactivating activity of ribozymes in animal cells.

Whilst the effectiveness of the ribozymes in vivo, is believed to be caused by one or more catalytic regions which are capable of cleavage of a target RNA, the presence of such regions in "Antisense" RNA type ribozymes may not actually lead to cleavage in vivo if the entire RNA/antisense RNA molecule does not fall apart. However, regardless of whether the molecule falls apart or not, the foregoing Examples demonstrate the effectiveness of the ribozyme in inactivating the target RNA. Thus, the invention is applicable to all ribozymes having a catalytic region capable of causing cleavage and a hybridizing region, regardless as to whether cleavage

41

actually occurs in the target RNA. i.e. The hybridizing region may be so large as to cause the combined RNA/ribozyme to stay together and prevent the target RNA being cleaved into separated
5 components even though the catalytic region is itself capable of causing cleavage.

10

CLAIMS:


1. A ribozyme comprising a hybridizing region which is complementary in nucleotide sequence to at least part of a target RNA, and a catalytic region which is adapted to cleave said target RNA wherein the hybridizing region contains 9 or more nucleotides.

2. A ribozyme comprising one or more arms formed of single stranded RNA and having a sequence complementary to at least part of a target RNA, said one or more arms being associated with a catalytic region, capable of cleaving said target RNA; and where the hybridizing region comprises a single arm of RNA said arm contains at least 9 nucleotides, and where the hybridizing region comprises 2 or more arms of RNA, the sum of nucleotides in said arms is greater than 9 nucleotides.

3. A ribozyme according to claim 2, having two single stranded arms of RNA, associated with one another through said catalytic region.

4. A ribozyme according to any preceding claim which does not modify cleaved target RNA.


5. A ribozyme of the formula:



wherein; X represents any ribonucleotide and each X residue may be the same or different;
the sum of n and n' is greater than 6 and n and n' may be the same or different;
an (*) represents a base pair between complementary ribonucleotides;
X' and X'' represent ribonucleotides of complementary sequence along at least part of their length to allow base pairing between the oligoribonucleotides, or X' and X'' together form a single RNA sequence wherein at least part of said sequence comprises a stem formed by base pairing between complementary nucleotides;
and optionally, an additional nucleotide selected from any one of A, G, C or U is inserted after ¹A.

6. A ribozyme as claimed in any preceding claim having the formula:

44



wherein; X represents any ribonucleotide and each X residue may be the same or different;
 an (*) represents a base pair between complementary ribonucleotides;
 n and n' are as previously defined;
 B represents a bond, a base pair, a ribonucleotide, or an oligoribonucleotide containing at least 2 ribonucleotides;
 m and m' are 1 or more and may be the same or different; and
 optionally, an additional nucleotide selected from any one of A, G, C or U is inserted after ¹A.

7. A ribozyme according to any one of claims 5 or 6, where the sum of n and n' is 16 or greater.

8. A ribozyme according to any preceding claim wherein the catalytic region is capable of cleaving a target RNA after the sequence X^oUY where X^o is any ribonucleotide, U is uracil and Y is adenine, cytosine or uracil.

9. A ribozyme containing multiple catalytic domains connected through flanking sequences, each said flanking sequence including sequences complementary to the target RNA which is desired to be inactivated.

10. A ribozyme according to any preceding claim which has been stabilized by circularization.

11. A ribozyme comprised of catalytic antisense sequences according to any preceding claim.

12. A method for the inactivation or cleavage of a target RNA which comprises reacting said target RNA with a ribozyme as claimed in any one of claims 1 to 11.

13. A method according to claim 12, wherein said inactivation occurs in-vivo in prokaryotic or eukaryotic cells, as a result of transcription of said ribozyme from a DNA or RNA transfer vector transfected into said cells or precursors thereof, or expression of a gene encoding said ribozyme which is integrated into the host genome, such that said ribozyme hybridizes to and inactivates a desired cellular target RNA.

14. A method according to claim 13, wherein said cells are plant or animal cells.

15. A method according to claim 13 or 14, wherein the target RNA is endogenous to the cells, or is exogenous, such as viral RNA.

16. A method according to claim 15, wherein target viral RNA is selected from RNA of the HIV(AIDS), hepatitis or other virus causing disease in mammals.

17. A method according to claim 15, wherein the viral RNA is bacteriophage RNA.

18. A method for the treatment of viral disease in man or animals which comprises administering to a subject in need of such treatment, an effective amount of a ribozyme according to any one of claims 1 to 11, said ribozyme hybridizing to and inactivating transcripts of said viral RNA.

19. A method for treating viral or viroid disease in plants which comprises administering to a plant a ribozyme according to any one of claims 1 to 11, said ribozyme hybridizing to and inactivating transcripts of said viral RNA.

20. A composition which comprises a ribozyme as claimed in any one of claims 1 to 11 in association with pharmaceutically, veterinarily, or agriculturally acceptable carrier or excipient.

21. A method for producing a ribozyme according to any one of claims 1 to 11 which comprises the steps of:

- (a) ligating into a transfer vector comprised of DNA, RNA or a combination thereof a nucleotide sequence corresponding to said ribozyme;
- (b) transcribing the nucleotide sequence of step (a) with RNA polymerase; and

(c) optionally purifying the ribozyme.

22. A transfer vector comprised of RNA or DNA or a combination thereof containing a nucleotide sequence which on transcription gives rise to a ribozyme as claimed in any one of claims 1 to 11.

23. A transfer vector according to claim 22, which is a bacterial plasmid or phage DNA.

24. A prokaryotic or eukaryotic cell containing nucleotide sequences which on transcription thereof give rise to a ribozyme as claimed in any one of claims 1 to 11.

2/19




Fig.2a.




Fig.2b.

3/19




Fig. 3.

4/19




Fig. 4a.

RzCAT-1 SITE:

#24




Fig. 4b.

5/19

RzCAT-2 SITE:




Fig. 4c.

RzCAT-3 SITE:




Fig. 4d.

6/19





Fig. 5a.

7/19

*Fig.5b.*

8/19




Fig. 6.

SUBSTITUTE SHEET

9/19






FIG. 7.

10/19






Fig.7. (cont.)

—●— SUBSTRATE —+— 5' PRODUCT

11/19





Fig. 8a.

12/19




FIG. 8a. (cont.)

14/19




Fig. 10a.




Fig. 10b.

15/19

RIBOZYME CEV9x(+) 5' GAAGUCCUUCAG 3'
 RIBOZYME CEV9x(-) 5' AGGGUCAGGUGA 3'
 RIBOZYME CEV25x(+) 5' GAAGUCGAGGUC 3'

Fig. 11.





Fig. 12b.

16/19




Fig. 13a.




Fig. 13b.

17/19




Fig. 14a.




Fig. 14b.

18/19




Fig.15a.




Fig.15b.




FIG. 16.




FIG. 17

SUBSTITUTE SHEET

INTERNATIONAL SEARCH REPORT

International Application No PCT/AU 88/00478

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) * According to International Patent Classification (IPC) or to both National Classification and IPC Int. Cl. ⁴ C12N 9/22, 5/02, 15/00, 1/20, C12P 19/34, C07H 21/02, A61K 31/73, 39/235, 39/285																							
II. FIELDS SEARCHED <div style="text-align: center; border-top: 1px solid black; border-bottom: 1px solid black;">Minimum Documentation Searched ⁷</div> <table style="width: 100%; border-collapse: collapse;"> <tr> <th style="width: 20%; text-align: left; border-bottom: 1px solid black;">Classification System</th> <th style="text-align: left; border-bottom: 1px solid black;">Classification Symbols</th> </tr> <tr> <td style="vertical-align: top; border-right: 1px solid black; padding: 5px;">IPC</td> <td style="padding: 5px;">WPI, WPIL, USPA DERWENT database keywords: RNA ENZYME; RIBOZYME; ENDORIBONUCLEASE and HYBRIDIZE</td> </tr> </table> <div style="text-align: center; border-top: 1px solid black; border-bottom: 1px solid black;">Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸</div>			Classification System	Classification Symbols	IPC	WPI, WPIL, USPA DERWENT database keywords: RNA ENZYME; RIBOZYME; ENDORIBONUCLEASE and HYBRIDIZE																	
Classification System	Classification Symbols																						
IPC	WPI, WPIL, USPA DERWENT database keywords: RNA ENZYME; RIBOZYME; ENDORIBONUCLEASE and HYBRIDIZE																						
AU: C12N 9/22, C07H 21/02, C12N 15/00, Chemical Abstracts Keywords as above																							
III. DOCUMENTS CONSIDERED TO BE RELEVANT ⁹ <table style="width: 100%; border-collapse: collapse;"> <tr> <th style="width: 10%; text-align: left; border-bottom: 1px solid black;">Category ⁹</th> <th style="text-align: left; border-bottom: 1px solid black;">Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²</th> <th style="text-align: left; border-bottom: 1px solid black;">Relevant to Claim No. ¹³</th> </tr> <tr> <td style="vertical-align: top; border-right: 1px solid black; padding: 5px;">X</td> <td style="padding: 5px;">Nature, volume 328, 13 August 1987, pp 596-600, Olke C. Uhlenbeck, "A small catalytic</td> <td style="vertical-align: top; padding: 5px;">(1,2,3,8,12, 22-24)</td> </tr> <tr> <td style="vertical-align: top; border-right: 1px solid black; padding: 5px;">Y</td> <td style="padding: 5px;">oligoribonucleotide"</td> <td style="vertical-align: top; padding: 5px;">(5,6)</td> </tr> <tr> <td style="vertical-align: top; border-right: 1px solid black; padding: 5px;">P,X P,Y</td> <td style="padding: 5px;">Nature, vol 334, number 6183, 18 August 1988, pp 585-591, J. Haseloff and W.L. Gerlach, "Simple RNA enzyme with new and highly specific endoribonuclease activities"</td> <td style="vertical-align: top; padding: 5px;">(1-13,22-24) (14-17)</td> </tr> <tr> <td style="vertical-align: top; border-right: 1px solid black; padding: 5px;">X</td> <td style="padding: 5px;">FEBS LETTERS, volume 220, number 2, August 1987, pp 283-287, Siegfried Boehm, "Similarities between a predicted secondary structure for the M₁ RNA ribozyme and the tRNA binding center of 16S rRNA from <u>E. coli</u>"</td> <td style="vertical-align: top; padding: 5px;">(1,2)</td> </tr> <tr> <td style="vertical-align: top; border-right: 1px solid black; padding: 5px;">P,X P,Y</td> <td style="padding: 5px;">Cold Spring Harbor Symposia on Quantitative Biology vol 41, Cold Spring Harbor Laboratory, 1987, pp 267-268, 274-275, J.R. Sampson et al, "Characterization of two RNA catalyzed cleavage reactions"</td> <td style="vertical-align: top; padding: 5px;">(1-3,8,12, 22-24) (5,6)</td> </tr> <tr> <td colspan="3" style="text-align: center; padding: 10px;">(continued)</td> </tr> </table>			Category ⁹	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³	X	Nature, volume 328, 13 August 1987, pp 596-600, Olke C. Uhlenbeck, "A small catalytic	(1,2,3,8,12, 22-24)	Y	oligoribonucleotide"	(5,6)	P,X P,Y	Nature, vol 334, number 6183, 18 August 1988, pp 585-591, J. Haseloff and W.L. Gerlach, "Simple RNA enzyme with new and highly specific endoribonuclease activities"	(1-13,22-24) (14-17)	X	FEBS LETTERS, volume 220, number 2, August 1987, pp 283-287, Siegfried Boehm, "Similarities between a predicted secondary structure for the M ₁ RNA ribozyme and the tRNA binding center of 16S rRNA from <u>E. coli</u> "	(1,2)	P,X P,Y	Cold Spring Harbor Symposia on Quantitative Biology vol 41, Cold Spring Harbor Laboratory, 1987, pp 267-268, 274-275, J.R. Sampson et al, "Characterization of two RNA catalyzed cleavage reactions"	(1-3,8,12, 22-24) (5,6)	(continued)		
Category ⁹	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³																					
X	Nature, volume 328, 13 August 1987, pp 596-600, Olke C. Uhlenbeck, "A small catalytic	(1,2,3,8,12, 22-24)																					
Y	oligoribonucleotide"	(5,6)																					
P,X P,Y	Nature, vol 334, number 6183, 18 August 1988, pp 585-591, J. Haseloff and W.L. Gerlach, "Simple RNA enzyme with new and highly specific endoribonuclease activities"	(1-13,22-24) (14-17)																					
X	FEBS LETTERS, volume 220, number 2, August 1987, pp 283-287, Siegfried Boehm, "Similarities between a predicted secondary structure for the M ₁ RNA ribozyme and the tRNA binding center of 16S rRNA from <u>E. coli</u> "	(1,2)																					
P,X P,Y	Cold Spring Harbor Symposia on Quantitative Biology vol 41, Cold Spring Harbor Laboratory, 1987, pp 267-268, 274-275, J.R. Sampson et al, "Characterization of two RNA catalyzed cleavage reactions"	(1-3,8,12, 22-24) (5,6)																					
(continued)																							
<div style="display: flex; justify-content: space-between;"> <div style="width: 48%;"> <p>* Special categories of cited documents: ¹⁰</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 48%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"Z" document member of the same patent family</p> </div> </div>																							
IV. CERTIFICATION <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%; border-bottom: 1px solid black; padding: 5px;"> Date of the Actual Completion of the International Search 23 March 1989 (23.03.89) </td> <td style="width: 50%; border-bottom: 1px solid black; padding: 5px;"> Date of Mailing of this International Search Report 30 MARCH 1989 (30.03.89) </td> </tr> <tr> <td style="border-bottom: 1px solid black; padding: 5px;"> International Searching Authority Australian Patent Office </td> <td style="border-bottom: 1px solid black; padding: 5px;"> Signature of Authorized Officer <div style="text-align: center;"> J.H. CHAN </div> </td> </tr> </table>			Date of the Actual Completion of the International Search 23 March 1989 (23.03.89)	Date of Mailing of this International Search Report 30 MARCH 1989 (30.03.89)	International Searching Authority Australian Patent Office	Signature of Authorized Officer <div style="text-align: center;"> J.H. CHAN </div>																	
Date of the Actual Completion of the International Search 23 March 1989 (23.03.89)	Date of Mailing of this International Search Report 30 MARCH 1989 (30.03.89)																						
International Searching Authority Australian Patent Office	Signature of Authorized Officer <div style="text-align: center;"> J.H. CHAN </div>																						

FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

P,X	FEBS LETTERS, volume 228, number 2, February 1988,	(1,2,3,8,12)
P,Y	pp 228-30, Koizumi, M., et al, "Construction of a series of several self-cleaving RNA duplexes using synthetic 21 mers"	(5,6)
A	Nature, vol 324(4), 4 December 1986, pp 429-433, Zaug et al, "The Tetrahymena ribozyme acts like an RNA restriction endonuclease".	(1,2,12)
A	Science, vol 231, 31 January 1986, pp 473-474, Zaug et al, "The intervening sequence RNA of Tetrahymena is an enzyme"	(1,12)
A	Science, vol 236, 19 June 1987, pp 1532-1539, T. Cech, "The Chemistry of Self-Splicing RNA enzymes"	(1,12)

V. ☒ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE ¹

This International search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. ☒ Claim numbers 18 because they relate to subject matter not required to be searched by this Authority, namely:
methods for treatment of the human or animal body by therapy (Rule 39 iv).
2. ☒ Claim numbers 1-4 because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
with respect to the phrase "a catalytic region" in the claims.
3. ☐ Claim numbers _____ because they are dependent claims and are not drafted in accordance with the second and third sentences of PCT Rule 6.4(a).

VI. ☐ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING ²

This International Searching Authority found multiple inventions in this international application as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.
2. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:
3. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:
4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the International Searching Authority did not invite payment of any additional fee.

Remark on Protest

- ☐ The additional search fees were accompanied by applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.